

Dissociating limit in density functional theory with Coulomb optimal transport cost

GUY BOUCHITTÉ, GIUSEPPE BUTTAZZO, THIERRY CHAMPION
AND LUIGI DE PASCALE

Abstract. In the framework of Density Functional Theory with Strongly Correlated Electrons we consider the so-called bond dissociating limit for the energy of an aggregate of atoms. We show that the multi-marginals optimal transport cost with Coulombian electron-electron repulsion describes a dissociation effect. The variational limit is completely calculated in the case of $N = 2$ electrons. The theme of fractional number of electrons appears naturally and brings into play the question of optimal partial transport cost. A plan is outlined to complete the analysis which involves the study of the relaxation of optimal transport cost with respect to the weak* convergence of measures.

Mathematics Subject Classification (2020): 82M30 (primary); 49J45, 49N15, 49K30 (secondary).

1. Introduction

The goal of this paper is to describe the behavior, as $\varepsilon \rightarrow 0$, of the asymptotically minimizing sequences $\{\rho_\varepsilon\}_\varepsilon$ of the functionals

$$F_\varepsilon(\rho) = \varepsilon T(\rho) + bC(\rho) - U(\rho), \quad (1.1)$$

where ε is a small parameter depending on the Planck constant \hbar , b is a given positive constant, and the three terms appearing are defined on the set of Borel probabilities over \mathbb{R}^3 as follows:

The research of the second and fourth authors is part of the project 2010A2TFX2 *Calcolo delle Variazioni* funded by the Italian Ministry of Research and is partially financed by the “*Fondi di ricerca di ateneo*” of the University of Firenze. The second and fourth authors are member of the Gruppo Nazionale per l’Analisi Matematica, la Probabilità e le loro Applicazioni (GNAMPA) of the Istituto Nazionale di Alta Matematica (INdAM).

Received March 25, 2019; accepted in revised form May 23, 2020.

Published online September 2021.

- The von Weizsäcker *kinetic energy* T is of the form

$$T(\rho) = \int \frac{|\nabla \rho|^2}{4\rho} dx = \int |\nabla \sqrt{\rho}|^2 dx ,$$

where we use the convention that integrals with no precised domain are taken over the whole ambient space \mathbb{R}^3 ;

- The *correlation term* C is given by means of the multi-marginal mass transport functional

$$C(\rho) := \inf \left\{ \int_{\mathbb{R}^{3N}} c(x_1, \dots, x_N) dP(x_1, \dots, x_N) : \forall i = 1, \dots, N, \pi_i^\# P = \rho \right\}, \quad (1.2)$$

where P is a probability on \mathbb{R}^{3N} , π_i is the projection map from \mathbb{R}^{3N} on its i -th factor \mathbb{R}^3 , $\#$ denotes the push forward operator defined for a map f and a measure μ by

$$f^\# \mu(E) = \mu(f^{-1}(E)),$$

and c is the Coulomb correlation function

$$c(x_1, \dots, x_N) = \sum_{1 \leq i < j \leq N} \frac{1}{|x_i - x_j|} .$$

Since c is symmetric, we may indifferently take the competitors P symmetric or not in (1.2). As it will be explained later, from the physical point of view $P \simeq |\psi|^2$ where ψ is a wave function, should satisfy the symmetry assumption. Nevertheless it will be more convenient to allow for non-symmetric competitors P along some proofs;

- The *potential term* U is of the form

$$U(\rho) = \int V(x) d\rho(x)$$

being $V(x)$ the Coulomb potential

$$V(x) = \sum_{1 \leq k \leq M} \frac{Z_k}{|x - X_k|} .$$

We shall also write U in the form

$$U(\rho) = \sum_{k=1}^M Z_k U_{X_k}(\rho) \quad \text{where} \quad U_{X_k}(\rho) := \int \frac{1}{|x - X_k|} d\rho(x) .$$

The Density Functional Theory (DFT) was introduced in the works of Thomas [25] and Fermi [10] and then revived by Hohenberg, Kohn and Sham [11, 12] and, from

a variational point of view, by Levy [14] and Lieb [15]. The aim of DFT is to develop a theory of Quantum Mechanics using the N -points probability distribution of electrons ρ (also known as charge density) as the main variable, instead of the wave functions. The usual choice for the definition of the electrons density is $\tilde{\rho} = N\rho$ so that, given a subset B of \mathbb{R}^3 , the quantity $\tilde{\rho}(B)$ represents the number of electrons which are in B . In this paper it will be more convenient to deal with probability densities. The functional F_ε above is a typical density functional modeling an aggregate of atoms. The correlation term represents the electron-electron interaction while the potential term represents the interaction of the electrons with the nuclei located in position X_k and with positive charge Z_k . More details on this model are given in the Appendix.

In [15, Theorems 1.1 and 1.2] Lieb proved that the physically appropriate space to consider as domain of the functional F_ε is

$$\mathcal{H} = \left\{ \rho \in L^1(\mathbb{R}^3) : \rho \geq 0, \int \rho dx = 1, \sqrt{\rho} \in H^1(\mathbb{R}^3) \right\}.$$

When the Coulomb correlation term C above is chosen, the theory is usually called Strongly Correlated Electrons Density Functional Theory (SCE-DFT); it was started and developed since the late 90's (see for instance [21–23]) and the connection with optimal transport was made in [4, 7].

We will show that, as $\varepsilon \rightarrow 0$, the minimal values of F_ε tend to $-\infty$ as $-1/\varepsilon$, so it will be convenient to consider the rescaled functionals

$$G_\varepsilon(\rho) = \varepsilon F_\varepsilon(\rho) = \varepsilon^2 T(\rho) + \varepsilon b C(\rho) - \varepsilon U(\rho) \quad (1.3)$$

that have the same minimizers as F_ε (we refer to Definition 3.4 for a precise statement). It turns out that, by a scaling property of the above functionals, minimizing G_ε is equivalent to minimizing the functional F_1 with nuclei at positions $\tilde{X}_i = X_i/\varepsilon$ (see (3.2)): in this respect the asymptotic study of G_ε corresponds to the study of the so-called *dissociating bond problem* (see in particular Remark 5.11 and [5, 18]). The goal is then to characterize the Γ -limit G of G_ε with respect to the weak* convergence of measures. In this way, the minimizers ρ_ε of G_ε (or equivalently of F_ε) will tend to minimizers of G in the weak* convergence of measures. Since we are on the whole space \mathbb{R}^3 , the weak* convergence in principle does not preserve the total mass, so we could expect that the limits ρ of ρ_ε are not anymore probabilities and only satisfy the inequality $\int d\rho \leq 1$.

The analysis of the limit functional G is then very important and the ultimate goal is to characterize G explicitly in terms of the data. This would allow to determine the measures ρ that minimize G and, by consequence, a precise picture of the shape of the the minimizers ρ_ε of F_ε as ε is small. In this respect, an interesting issue is to establish whether or not limiting measures ρ which are minimal for G are still probability measures. In view of the scaling above, we expect that, for a given distribution of nuclei $\{X_k, Z_k\}$, $G(\rho)$ depends only on the weights $\alpha_k := \rho(\{X_k\})$ of ρ on each nuclei, namely:

$$G(\rho) = \gamma(Z_1, \dots, Z_M, \alpha_1, \dots, \alpha_M) = \gamma(Z, \alpha),$$

being $\gamma(Z, \cdot)$ a function defined on the simplex

$$\left\{ \alpha \in [0, 1]^M : \sum_{k=1}^M \alpha_k \leq 1 \right\}.$$

Then showing that $\gamma(Z, \cdot)$ achieves its minimum at α such that $\sum \alpha_k = 1$ would imply that the probability $\rho = \sum_k \alpha_k \delta_{X_k}$ is optimal for G . Note that in this case, the uniqueness of such α would imply as a byproduct the tight convergence $\rho_\varepsilon \rightarrow \rho$ for any minimizing sequence $\{\rho_\varepsilon\}$.

In principle the function γ above could be very involved, mixing the data in a very intricate way; in fact a better situation occurs if

$$\gamma(Z_1, \dots, Z_M, \alpha_1, \dots, \alpha_M) = \sum_{k=1}^M g(Z_k, \alpha_k) \quad (1.4)$$

where the function g can be deduced through the solution of an *auxiliary problem* with only one nucleus. Several points on the asymptotic analysis of the functionals F_ε can be achieved in full generality. However, due to some technical difficulties, mostly related to the analysis of the correlation term C , we are able to obtain a complete characterization of the Γ -limit functional G only when the number N of electrons is at most 2 (for any number M of nuclei). In this case, we are able to obtain G explicitly by means of a notion of *fractional transport cost*. This will imply that the optimal measures ρ for G are actually probabilities, and so the convergence of ρ_ε to ρ is in the narrow sense. In addition (see Theorem 5.7), formula (1.4) is shown to hold, together with an expression of the function g that can be deduced by means of an auxiliary variational problem.

The plan of the paper is the following. In Sections 2 and 3 we introduce the notation used in the following and show some basic properties of the functionals F_ε and of its three components. We also introduce the functionals G_ε together with some of their asymptotic properties. Section 4 is devoted to the non-interacting case $b = 0$, in which the electron-electron correlation term is not present. This simplifies a lot the analysis and an explicit characterization of the Γ -limit functional is obtained in this case. In Section 5 we treat the case with the electron-electron interaction term $C(\rho)$ and we provide, in the case $N = 2$, a general expression of the Γ -limit functional G . Finally, in the last section of the paper we discuss about the case $N > 2$ and some open issues, concluding the paper with some comments about our future work program. A short appendix contains some background on the links between DFT and the Born-Oppenheimer model for the electronic structure of several particles systems, including the bond dissociating limit for that model.

ACKNOWLEDGEMENTS. This paper has been written during some visits of the authors at the Department of Mathematics of University of Firenze, University of Pisa, and at the Laboratoire IMATH of University of Toulon. The last part of the work

was completed during the visiting fellowship of the first author to CRM-CNRS Laboratorio Fibonacci in Pisa. The authors gratefully acknowledge the warm hospitality and support of these institutions. During the work, several conversations with Simone Di Marino, Paola Gori-Giorgi, Mathieu Lewin and Michael Seidl helped to understand the physical problem and to better frame it in the literature; they are all warmly acknowledged.

2. Introductory results on the correlation term C

In the following by \mathcal{P} we denote the class of Borel probabilities in \mathbb{R}^3 and by \mathcal{P}^- the class of Borel subprobabilities on \mathbb{R}^3 , that is Borel measures μ with $\int d\mu \leq 1$. By $\xrightarrow{*}$ and \rightarrow we respectively denote the weak* convergence and the narrow convergence on \mathcal{P} and on \mathcal{P}^- , related to the duality with continuous compactly supported functions and with bounded continuous functions. Finally, we indicate by δ_X the Dirac mass at the point X and by $\|\rho\|$ the quantity $\int d\rho$.

By $T(\rho), C(\rho), U(\rho)$ we denote the functionals detailed in the Introduction, representing respectively the kinetic energy, the correlation term, and the potential term of a density $\rho \in \mathcal{P}$. As noted in the introduction, the absolutely minimizing sequences for the functionals F_ε (or G_ε) may only weak* converge to elements in \mathcal{P}^- , it is then convenient to extend the functionals $T(\rho), C(\rho)$ and $U(\rho)$ to any non-negative bounded measure (in particular on \mathcal{P}^-) by 1-homogeneity. We shall denote these 1 homogeneous extensions by $\mathcal{T}(\rho), \mathcal{C}(\rho)$ and $\mathcal{U}(\rho)$ respectively: for \mathcal{T} and \mathcal{U} , this extension process obviously leads to the same expression, and these two functionals are lower semicontinuous on \mathcal{P}^- for the weak* convergence. For the correlation term $C(\rho)$, we also define a 1-homogenous extension by setting $C(\rho) := \|\rho\| C(\frac{\rho}{\|\rho\|})$ which can be rewritten in a similar way as in (1.2), namely

$$C(\rho) = \inf \left\{ \int_{\mathbb{R}^{3N}} c(x_1, \dots, x_N) dP(x_1, \dots, x_N) : \pi_i^\# P = \rho, \forall i = 1, \dots, N \right\} \quad (2.1)$$

where now transport plans P are non-negative Borel measures on R^{3N} with total mass $\|P\| = \|\rho\|$.

While most of the readers are probably familiar with the first and third terms T and U of F_ε , we believe that the following results are useful to understand the correlation term C .

Proposition 2.1. *For every probability ρ on \mathbb{R}^3 we have*

$$C(\rho) \leq \frac{N(N-1)}{2} \int_{(\mathbb{R}^3)^2} \frac{1}{|x-y|} d\rho(x)d\rho(y).$$

In particular, $C(\rho)$ is finite for every probability density ρ which is bounded and with compact support.

Proof. From the definition of $C(\rho)$, we may take $\rho \otimes \rho \otimes \dots \otimes \rho$ as a particular probability P , so that

$$C(\rho) \leq \int_{\mathbb{R}^{3N}} c(x_1, \dots, x_N) d\rho(x_1) \dots d\rho(x_N)$$

and this last integral reduces to

$$\frac{N(N-1)}{2} \int_{(\mathbb{R}^3)^2} \frac{1}{|x-y|} d\rho(x) d\rho(y)$$

by the symmetry of the function c . □

Proposition 2.2. *For every probability ρ on \mathbb{R}^3 we have*

$$C(\rho) \geq \frac{N(N-1)}{4\sqrt{\text{Var}(\rho)}},$$

where $\text{Var}(\rho)$ denotes the variance of the probability ρ . In particular, when ρ is a Dirac mass, then $\text{Var}(\rho) = 0$ and we recover that $C(\rho) = +\infty$.

Proof. Let P be a probability on \mathbb{R}^{3N} with $\pi_i^\# P = \rho$ for all $i = 1, \dots, N$ and let $\gamma = (\pi_i \times \pi_j)^\# P$ be the projection of P on the product $\mathbb{R}^3 \times \mathbb{R}^3$. Then we have

$$1 \leq \left(\int_{(\mathbb{R}^3)^2} \frac{1}{\sqrt{|x_i - x_j|}} \sqrt{|x_i - x_j|} d\gamma \right)^2 \leq \int_{(\mathbb{R}^3)^2} \frac{1}{|x_i - x_j|} d\gamma \int_{(\mathbb{R}^3)^2} |x_i - x_j| d\gamma.$$

Without loss of generality we assume $\text{Var}(\rho) < +\infty$ (otherwise there is nothing to prove), so that ρ has a finite expectation $\mathbb{E}(\rho) = \int x d\rho$ and we can write

$$\begin{aligned} \int_{(\mathbb{R}^3)^2} |x_i - x_j| d\gamma &\leq \int_{(\mathbb{R}^3)^2} (|x_i - \mathbb{E}(\rho)| + |\mathbb{E}(\rho) - x_j|) d\gamma \\ &\leq 2 \int |x - \mathbb{E}(\rho)| d\rho(x) \leq 2\sqrt{\text{Var}(\rho)}. \end{aligned}$$

Thus

$$\int_{(\mathbb{R}^3)^2} \frac{1}{|x_i - x_j|} d\gamma \geq \frac{1}{2\sqrt{\text{Var}(\rho)}}.$$

Summing on all $1 \leq i < j \leq N$ we obtain

$$\int_{\mathbb{R}^{3N}} c(x_1, \dots, x_N) dP(x_1, \dots, x_N) \geq \frac{N(N-1)}{4\sqrt{\text{Var}(\rho)}}$$

as required. □

Remark 2.3. By 1-homogeneity, Propositions 2.1 and 2.2 yield

$$\forall \rho \in \mathcal{P}^-, \quad \frac{N(N-1) \|\rho\|^2}{4\sqrt{\text{Var}(\rho)}} \leq \mathcal{C}(\rho) \leq \frac{N(N-1)}{2 \|\rho\|} \int_{(\mathbb{R}^3)^2} \frac{1}{|x-y|} d\rho(x)d\rho(y).$$

In particular $0 < \mathcal{C}(\rho)$ for any non zero smooth $\rho \in \mathcal{P}^-$ with compact support.

It turns out that the function $\mathcal{C}(\rho)$ is lower semicontinuous with respect to the tight convergence of probability measures. However this is not sufficient for our purposes since sequences with uniformly bounded energy (1.3) are not tight in general. Therefore we will deal with the weak* convergence and accordingly it is useful to introduce the following lower semicontinuous extension of \mathcal{C} to subprobabilities:

$$\forall \rho \in \mathcal{P}^-, \quad \overline{\mathcal{C}}(\rho) := \inf \left\{ \liminf_{n \rightarrow +\infty} \mathcal{C}(\rho_n) : \rho_n \in \mathcal{P}, \rho_n \xrightarrow{*} \rho \right\}.$$

A natural guess could be that $\overline{\mathcal{C}}(\rho)$ is equal to $\mathcal{C}(\rho)$; however, this is not the case as the following proposition shows.

Proposition 2.4. *The functional \mathcal{C} is convex on \mathcal{P}^- , and the lower semicontinuous envelope $\overline{\mathcal{C}}$ with respect to the weak* convergence satisfies*

$$\overline{\mathcal{C}}(\rho) \leq \min \left\{ \mathcal{C}(\mu) : 0 \leq \mu \leq \rho, \int d\mu \geq \frac{N}{N-1} \left[\int d\rho - \frac{1}{N} \right] \right\}. \quad (2.2)$$

Note that as a corollary of (2.2) it holds

$$\int d\rho \leq \frac{1}{N} \implies \overline{\mathcal{C}}(\rho) = 0 \quad \forall \rho \in \mathcal{P}^-$$

which together with Remark 2.3 yields that $\overline{\mathcal{C}} \neq \mathcal{C}$.

Proof. The convexity of \mathcal{C} follows from the linearity of the constraint $\pi_i^\# P = \rho$ for all i .

For the second statement, we first note that when $\int d\rho \leq 1/N$, the minimum in (2.2) is clearly 0 and attained for $\mu = 0$. Moreover, since $\overline{\mathcal{C}}$ is 1-homogeneous the second constraint in (2.2) may be replaced by $\int d\mu = \frac{N}{N-1} \left[\int d\rho - \frac{1}{N} \right]$ whenever $\int d\rho \geq 1/N$. Note that this minimum is then attained since the set of measures

$$\left\{ \mu : 0 \leq \mu \leq \rho, \int d\mu = \frac{N}{N-1} \left[\int d\rho - \frac{1}{N} \right] \right\}$$

is tight and \mathcal{C} is lower semicontinuous for the narrow convergence. We also note that (2.2) obviously holds when $\rho \in \mathcal{P}$ since $\mu = \rho$ is then the only admissible choice.

Now let $\rho \in \mathcal{P}^-$ with $\int d\rho < 1$, we first assume that $\int d\rho \geq 1/N$. Let μ such that $0 \leq \mu \leq \rho$, $\int d\mu = \frac{N}{N-1} \left[\int d\rho - \frac{1}{N} \right]$ and μ optimal in (2.2). We set $\nu = \rho - \mu$, so that $\int d\rho + (N - 1) \int d\nu = 1$. Let $\xi_1 = 0$, consider $N - 1$ distinct vectors $\xi_2, \dots, \xi_N \in \mathbb{R}^3 \setminus \{0\}$, and set for every $n \in \mathbb{N}$

$$\rho_{n,k} = \tau_{n,k} \# \nu, \quad k \in \{1, \dots, N\},$$

where $\tau_{n,k} := \tau_{(n \xi_k)}$, being τ_ξ the translation by ξ defined on \mathbb{R}^3 by $\tau_\xi : x \mapsto x + \xi$. We can now set

$$\forall n, \quad \rho_n := \rho + \sum_{k=2}^N \rho_{n,k} = \mu + \sum_{k=1}^N \rho_{n,k}$$

and we note that ρ_n belongs to \mathcal{P} for any n and $\rho_n \xrightarrow{*} \rho$ as $n \rightarrow +\infty$. We also denote by P an optimal plan for $\mathcal{C}(\mu)$, then we set

$$\forall n, \quad P_n := P + \sum_{i=1}^N (\tau_{n,\sigma^i(1)} \times \dots \times \tau_{n,\sigma^i(N)}) \# \nu,$$

where σ is the permutation on $\{1, \dots, N\}$ such that $\sigma(j) = j + 1$ and $\sigma(N) = 1$. Then for all n the plan P_n satisfies

$$\forall j \in \{1, \dots, N\}, \quad \pi_j \# P_n = \mu + \sum_{i=1}^N \tau_{n,\sigma^i(j)} \# \nu = \rho_n$$

so P_n is admissible for $\mathcal{C}(\rho_n)$. We can now estimate

$$\begin{aligned} \mathcal{C}(\rho_n) &\leq \int_{\mathbb{R}^{3N}} c dP_n = \int_{\mathbb{R}^{3N}} c dP + \sum_{i=1}^N \int c(\tau_{n,\sigma^i(1)}(x), \dots, \tau_{n,\sigma^i(N)}(x)) \nu(dx) \\ &= \int_{\mathbb{R}^{3N}} c dP + N \sum_{1 \leq i < j \leq N} \frac{\|\nu\|}{n |\xi_i - \xi_j|} \end{aligned}$$

from which we conclude that

$$\overline{\mathcal{C}}(\rho) \leq \liminf_{n \rightarrow +\infty} \mathcal{C}(\rho_n) \leq \int_{\mathbb{R}^{3N}} c dP = \mathcal{C}(\mu).$$

as required.

It remains to treat the case $\int d\rho \leq \frac{1}{N}$. It follows from the preceding that $\overline{\mathcal{C}}(\rho) = 0$ for any ρ such that $\int d\rho = \frac{1}{N}$, then by weak* lower semicontinuity of $\overline{\mathcal{C}}$ and by approximation this also holds for any $\rho \in \mathcal{P}^-$ with $\int d\rho \leq \frac{1}{N}$, which finishes the proof of (2.2) in this case. \square

The full characterization of \bar{C} on \mathcal{P}^- is a quite involved problem, in particular it is an open problem whether equality holds in (2.2) for $N \geq 3$ but holds true for $N = 2$.

Proposition 2.5. *In the case $N = 2$ it holds*

$$\bar{C}(\rho) = \min \left\{ \mathcal{C}(\mu) : 0 \leq \mu \leq \rho, \int d\mu \geq 2 \int d\rho - 1 \right\} \quad \forall \rho \in \mathcal{P}^-, \quad (2.3)$$

with $\mathcal{C}(\mu)$ defined in (2.1)

Proof. In order to obtain the reverse inequality of (2.2), consider a sequence ρ_n in \mathcal{P} weakly* converging to $\rho \in \mathcal{P}^-$. For each n we denote by P_n an optimal plan for ρ_n , then we may assume that $P_n \xrightarrow{*} P$ for some non-negative Borel measure P over $(\mathbb{R}^3)^2$, with marginals $\pi_1^\# P = \pi_2^\# P = \mu$ for some $\mu \in \mathcal{P}^-$. Then one gets

$$\begin{aligned} \liminf_{n \rightarrow +\infty} C(\rho_n) &= \liminf_{n \rightarrow +\infty} \int_{(\mathbb{R}^3)^2} c(x_1, x_2) P_n(dx_1, dx_2) \\ &\geq \int_{(\mathbb{R}^3)^2} c(x_1, x_2) P(dx_1, dx_2) \geq \mathcal{C}(\mu). \end{aligned}$$

We first claim that $\mu \leq \rho$: indeed, let $\phi, \psi \in C_c^\infty(\mathbb{R}^3)$ be non-negative and $0 \leq \psi \leq 1$, then we have

$$\begin{aligned} \langle \rho, \phi \rangle &= \lim_{n \rightarrow +\infty} \int_{(\mathbb{R}^3)^2} \phi(x) P_n(dx, dy) \\ &\geq \lim_{n \rightarrow +\infty} \int_{(\mathbb{R}^3)^2} \phi(x) \psi(y) P_n(dx, dy) \\ &= \int_{(\mathbb{R}^3)^2} \phi(x) \psi(y) P(dx, dy). \end{aligned}$$

Now letting $\psi \nearrow 1$ we obtain by the monotone convergence theorem that

$$\langle \rho, \phi \rangle \geq \int_{(\mathbb{R}^3)^2} \phi(x) P(dx, dy) = \langle \mu, \phi \rangle$$

and, since this is true for any non-negative test function ϕ , we get the claim.

It remains to prove that $\int d\mu \geq 2 \int d\rho - 1$. For this, consider a ball B_R centered at the origin such that $\rho(\partial B_R) = 0$; from what seen above this implies $\mu(\partial B_R) = P(\partial B_R^2) = 0$. Then for all $n \in \mathbb{N}$ one has

$$\rho_n(B_R) = P_n(B_R^2) + P_n(B_R \times B_R^c) \leq P_n(B_R^2) + \rho_n(B_R^c) = P_n(B_R^2) + 1 - \rho_n(B_R)$$

so that, passing to the limit as $n \rightarrow \infty$ gives

$$2\rho(B_R) - 1 \leq P(B_R^2) \leq \mu(B_R),$$

and then the claim follows by letting R go to $+\infty$. □

Remark 2.6. As a corollary of Propositions 2.4 and 2.5, it appears that when $N = 2$ one has $\overline{C}(\rho) > 0$ if and only if $\int d\rho \geq 1/2$: indeed, if $\int d\rho \geq 1/2$, since the minimum giving $\overline{C}(\rho)$ is attained for some μ with $\int d\mu = 2 \int d\rho - 1 > 0$, we obtain $\mu \neq 0$ so that $\overline{C}(\rho) = C(\mu) > 0$.

3. Basic inequalities, properties of the sequence and rescaling

We now get back to the preliminary analysis of F_ε . As proved by Lieb [15, Theorem 1.1 and Theorem 1.2] the correct space where to minimize the functional F_ε is

$$\mathcal{H} = \left\{ \rho \in L^1(\mathbb{R}^3) : \rho \geq 0, \int \rho dx = 1, \sqrt{\rho} \in H^1(\mathbb{R}^3) \right\}.$$

Proposition 3.1. *For every $\varepsilon > 0$ the functional F_ε is convex.*

Proof. To prove convexity we look separately at the three terms which compose the functional F_ε . The first term is convex with respect to the pair $(\rho, \nabla\rho)$ thanks to the convexity of the function $(s, v) \mapsto |v|^2/s$ on $\mathbb{R}^+ \times \mathbb{R}^d$. We notice that the kinetic energy can be also written in the form

$$T(\rho) = \int |\nabla\sqrt{\rho}|^2 dx.$$

The transport cost $C(\rho)$ defined in (1.2) is a linear problem with respect to the probability P on \mathbb{R}^{3N} , with linear constraints $\pi_i^\# P = \rho$, so that it is convex. The term $U(\rho)$ is linear with respect to ρ . □

The different behavior of the three terms with respect to the action of homotheties on the probability measures has a relevance in the study of the Γ -limit.

Definition 3.2. For every probability measure $\rho \in \mathcal{P}$ and every $s > 0$ we set

$$\rho^s = h_s^\# \rho$$

where $h_s : x \mapsto x/s$.

Remark 3.3. Note in particular that if ρ has a compact support then the support of ρ^s is $h_s(\text{supp}(\rho))$. We shall use in the following sections that $\rho^s \rightharpoonup \delta_0$ as $s \rightarrow +\infty$, which straightly follows from the definition. Also notice that the map $\rho \mapsto \rho^s$ has for inverse $\rho \mapsto \rho^{1/s}$. Finally, since the elements $\rho \in \mathcal{P}$ for which the functionals G_ε are defined are in L^1 , it is interesting to note that in this case the probability ρ^s is also in L^1 with density

$$\rho^s(x) = s^3 \rho(sx).$$

When $M = 1$ and $X_1 = 0$, a simple calculation shows that the three terms in F_ε scale as follows:

$$T(\rho^s) = s^2T(\rho), \quad C(\rho^s) = sC(\rho), \quad U(\rho^s) = U_0(\rho^s) = sU_0(\rho).$$

Therefore, minimizing the functional (1.1) with respect to ρ^s leads to minimize the quantity

$$\varepsilon s^2T(\rho) + s(bC(\rho) - U(\rho)).$$

A first minimization with respect to the variable $s \geq 0$ reduces the problem $\inf_\rho F_\varepsilon$ to minimize with respect to ρ the ratio

$$-\frac{1}{4\varepsilon} \frac{(U(\rho) - bC(\rho))_+^2}{T(\rho)}$$

where $(\cdot)_+$ denotes the positive part function. Then

$$\inf_{\rho \in \mathcal{P}} F_\varepsilon(\rho) = -\frac{K}{\varepsilon}, \tag{3.1}$$

where

$$K = \sup_{\rho \in \mathcal{P}} \frac{(U(\rho) - bC(\rho))_+^2}{4T(\rho)}.$$

This last supremum is finite as a consequence of the inequality $(U - bC)_+^2 \leq U^2$ and of Lemma 3.6 below applied to $u^2 = \rho$ for any probability ρ in the domain of F_ε .

When $M > 1$ the potential term U does not have anymore the scaling property above so we use the following estimate (setting $Z = \sum_{1 \leq k \leq M} Z_k$):

$$F_\varepsilon(\rho) = \sum_{1 \leq k \leq M} \frac{Z_k}{Z} \left[\varepsilon T(\rho) + bC(\rho) - \int \frac{Z\rho(x)}{|x - X_k|} dx \right],$$

so that

$$\begin{aligned} & \sum_{1 \leq k \leq M} \frac{Z_k}{Z} \inf_\rho \left[\varepsilon T(\rho) + bC(\rho) - \int \frac{Z\rho(x)}{|x - X_k|} dx \right] \leq \inf_\rho F_\varepsilon(\rho) \\ & \leq \inf_\rho \left[\varepsilon T(\rho) + bC(\rho) - \int \frac{Z_1\rho(x)}{|x - X_1|} dx \right]. \end{aligned}$$

It follows that, for suitable positive constants K_1 and K_2 , we have

$$-\frac{K_1}{\varepsilon} \leq \inf_\rho F_\varepsilon(\rho) \leq -\frac{K_2}{\varepsilon}.$$

In conclusion the minimal values of F_ε tend to $-\infty$ with order $1/\varepsilon$. This justifies the introduction of the rescaled functionals

$$G_\varepsilon(\rho) = \varepsilon F_\varepsilon(\rho).$$

Due to the lack of compactness for the tight convergence, it is not clear whether the functionals F_ε and G_ε admit minimizers in \mathcal{P} . As we are interested in the asymptotics as $\varepsilon \rightarrow 0$, we will use the following asymptotic notion of minimizer.

Definition 3.4. We say that $\{\rho_\varepsilon\}_\varepsilon \in \mathcal{P}$ is asymptotically minimizing for G_ε (respectively for F_ε) whenever

$$\lim_{\varepsilon \rightarrow 0} \left[G_\varepsilon(\rho_\varepsilon) - \inf_{\mathcal{P}} G_\varepsilon \right] = 0 \quad \left(\text{respectively } \lim_{\varepsilon \rightarrow 0} \frac{F_\varepsilon(\rho_\varepsilon) - \inf_{\mathcal{P}} F_\varepsilon}{\varepsilon} = 0 \right).$$

Let us emphasize that such a sequence of asymptotic minimizers converge (up to subsequences) weakly* to an element of \mathcal{P}^- . We may now apply to the family G_ε the Γ -convergence theory in order to identify the Γ -limit functional G and so, as a consequence, the behaviour of the asymptotically minimizing sequences, that will converge to minimizers of G .

The functional G_ε has also a physical interest by itself and deserves to be written explicitly

$$G_\varepsilon(\rho) = \varepsilon^2 T(\rho) + \varepsilon bC(\rho) - \varepsilon U(\rho).$$

The same homogeneities of the terms in the functional F_ε allow us to rewrite

$$G_\varepsilon(\rho) = T(\rho^\varepsilon) + bC(\rho^\varepsilon) - \sum_{1 \leq k \leq M} \int \frac{Z_k \rho^\varepsilon(x)}{|x - X_k/\varepsilon|} dx. \quad (3.2)$$

Then letting ε go to 0 is equivalent to let the distance between the nuclei go to $+\infty$. For this reason, when considering a molecule, the limit as $\varepsilon \rightarrow 0$ of G_ε models the dissociation of chemical bonds between the atoms composing the molecule.

In the following we denote by G^+ and by G^- respectively the Γ -limsup and the Γ -liminf of the family G_ε . Since the space \mathcal{P}^- endowed with the weak* convergence is metrizable and compact, by the general theory of the Γ -convergence (see for instance [9]) we have that a subsequence of G_ε (that we still continue to denote by G_ε) Γ -converges to some functional G . If we are able to fully characterize this limit functional G independently of the subsequence, we obtain that the full family G_ε is Γ -convergent to G . Therefore, in the following we may assume that G_ε Γ -converges to some functional G and we concentrate our efforts in obtaining a characterization of G in terms of the data only.

Since in general weak* limits of sequences of probabilities only belong to \mathcal{P}^- , we consider G^- , G^+ , and G as defined on \mathcal{P}^- . As a basic consequence of Γ -convergence theory (see [1]) we have the following result.

Proposition 3.5. *We have $G^- \leq G^+$; moreover the functionals G^- and G^+ are both weakly* lower-semicontinuous, and G^+ is convex.*

If we forget about the electron-electron interaction, *i.e.*, the optimal transport term C in G_ε , we obtain an estimate from below and at the same time an easier problem to work with. We will refer to this as the “*non-interacting case*” and the corresponding functionals will be denoted as

$$G_\varepsilon^0 = \varepsilon^2 T(\rho) - \varepsilon U(\rho).$$

We first characterize a wide space on which $G = G^+ = G^- = 0$.

Lemma 3.6. *There exists a constant κ such that for every domain Ω (bounded or not) we have*

$$\left[\int_\Omega \frac{u^2}{|x|} dx \right]^2 \leq \kappa \int_\Omega u^2 dx \int |\nabla u|^2 dx \quad \forall u \in H^1(\mathbb{R}^3).$$

Proof. Using the embedding of $H^1(\mathbb{R}^3)$ into $L^6(\mathbb{R}^3)$ we have for every $\delta > 0$

$$\begin{aligned} \int_\Omega \frac{u^2}{|x|} dx &\leq \int_{\Omega \setminus B_\delta} \frac{u^2}{\delta} dx + \int_{B_\delta} \frac{u^2}{|x|} dx \\ &\leq \frac{1}{\delta} \int_\Omega u^2 dx + \|u\|_{L^6(\mathbb{R}^3)}^2 \|1/|x|\|_{L^{3/2}(B_\delta)} \\ &\leq \frac{1}{\delta} \int_\Omega u^2 dx + \kappa \delta \int |\nabla u|^2 dx. \end{aligned}$$

Optimizing with respect to δ gives

$$\int_\Omega \frac{u^2}{|x|} dx \leq 2 \left[\kappa \int_\Omega u^2 dx \int |\nabla u|^2 dx \right]^{1/2}$$

as required. □

Proposition 3.7. *For every probability ρ we have*

$$-\frac{\kappa M}{4} \sum_{1 \leq k \leq M} Z_k^2 \rho(\{X_k\}) \leq G^-(\rho) \leq G^+(\rho) \leq 0, \tag{3.3}$$

where κ is given in Lemma 3.6. In particular $G^-(\rho) = G^+(\rho) = 0$ for every probability ρ that does not charge any of the points X_k .

Proof. By Proposition 2.1 we have for every smooth ρ with compact support

$$G_\varepsilon(\rho) = \varepsilon^2 T(\rho) + \varepsilon bC(\rho) - \varepsilon U(\rho) \leq K\varepsilon$$

for a suitable constant K depending on ρ . Therefore

$$G^+(\rho) \leq \lim_{\varepsilon \rightarrow 0} G_\varepsilon(\rho) \leq 0,$$

and the last inequality in (3.3) follows by approximation and by the lower semicontinuity of G^+ .

Let now ρ_ε be a generic sequence weakly* converging to ρ ; since the transport cost $C(\rho_\varepsilon)$ is nonnegative we have, setting $u_\varepsilon^2 = \rho_\varepsilon$,

$$G_\varepsilon(\rho_\varepsilon) \geq \varepsilon^2 \int |\nabla u_\varepsilon|^2 dx - \varepsilon \sum_{1 \leq k \leq M} Z_k \left[\int_{B_\delta(X_k)} \frac{u_\varepsilon^2}{|x - X_k|} dx + \int_{(B_\delta(X_k))^c} \frac{u_\varepsilon^2}{|x - X_k|} dx \right].$$

By using Lemma 3.6 and the fact that $|x - X_k| \geq \delta$ on $B_\delta(X_k)^c$ we obtain

$$G_\varepsilon(\rho_\varepsilon) \geq \varepsilon^2 \int |\nabla u_\varepsilon|^2 dx - \varepsilon \sum_{1 \leq k \leq M} Z_k \left[\kappa \int_{B_\delta(X_k)} u_\varepsilon^2 dx \int |\nabla u_\varepsilon|^2 dx \right]^{1/2} - \varepsilon \frac{Z}{\delta}.$$

Since $\varepsilon^2 A - \varepsilon B \geq -B^2/(4A)$, the sum of the first two terms in the last line gives

$$G_\varepsilon(\rho_\varepsilon) \geq -\frac{\kappa}{4} \left(\sum_{1 \leq k \leq M} Z_k \left[\int_{B_\delta(X_k)} u_\varepsilon^2 dx \right]^{1/2} \right)^2 - \varepsilon \frac{Z}{\delta}.$$

As $\varepsilon \rightarrow 0$ we obtain, for every $\delta > 0$,

$$G^-(\rho) \geq -\frac{\kappa}{4} \left(\sum_{1 \leq k \leq M} Z_k \left[\rho(\overline{B_\delta(X_k)}) \right]^{1/2} \right)^2,$$

and finally, as $\delta \rightarrow 0$,

$$G^-(\rho) \geq -\frac{\kappa}{4} \left(\sum_{1 \leq k \leq M} Z_k \left[\rho(\{X_k\}) \right]^{1/2} \right)^2 \geq -\frac{\kappa M}{4} \sum_{1 \leq k \leq M} Z_k^2 \rho(\{X_k\})$$

which concludes the proof. □

4. The non-interacting case

In this case, thanks to the absence of the transport term C , we are able to identify the limit functional G in a complete way. In order to stress the fact that $b = 0$ we denote the sequence by G_ε^0 and the limit by G^0 .

4.1. The hydrogen atom

The simplest case is $N = M = 1$; in other words we have a single nucleus with charge Z located at a point X_1 (that without loss of generality we can take the origin) and a single electron. In this situation the non interacting case maintains a physical meaning. Problem (3.1) then reduces to

$$\begin{aligned} \inf_{\rho \in \mathcal{P}} F_\varepsilon^0(\rho) &= \inf_{\rho \in \mathcal{P}} \{ \varepsilon T(\rho) - U(\rho) \} = \inf_{\rho \in \mathcal{P}} \left\{ -\frac{1}{4\varepsilon} \frac{(U(\rho))^2}{T(\rho)} \right\} \\ &= \frac{Z^2}{\varepsilon} \inf_{\rho \in \mathcal{P}} \left\{ -\frac{(U_0(\rho))^2}{4T(\rho)} \right\}. \end{aligned}$$

The value of the problem on the right-hand side is the first eigenvalue (negative) of the operator

$$-\Delta - \frac{1}{|x|}.$$

This is known to be equal to $-1/4$ with eigenfunctions proportional to

$$\bar{\rho} = \frac{1}{32\pi} e^{-|x|/2}$$

(see for instance [16, Example 11.10]). Summarizing, in the case $N = M = 1$ the minimizer $\bar{\rho}_\varepsilon$ of the functional F_ε^0 (or equivalently of the rescaled functional G_ε^0) is equal to

$$\bar{\rho}_\varepsilon(x) = \bar{\rho}^{1/\varepsilon}(x) = \frac{1}{4\pi(2\varepsilon)^3} e^{-|x|/(2\varepsilon)},$$

and tends, as $\varepsilon \rightarrow 0$, to the measure $\bar{\rho}(x) = \delta_0$ with the minimal values $G_\varepsilon(\bar{\rho}_\varepsilon) \rightarrow -Z^2/4$. In fact, by applying Theorem 4.4 to this particular case, we infer that the Γ -limit functional G^0 is identified on \mathcal{P}^- as

$$G^0(\rho) = -\frac{Z^2}{4} \rho(\{0\}).$$

From the previous discussion we deduce

$$-\frac{Z^2}{4} = \inf_{\rho \in \mathcal{P}} \{ T(\rho) - Z U_0(\rho) \} = \inf_{\rho} \left\{ \int \frac{|\nabla \rho|^2}{4\rho} dx - \int \frac{Z\rho}{|x|} dx \right\}. \quad (4.1)$$

4.2. The general case $N, M \geq 1$

We start by a localization lemma.

Lemma 4.1 (Localization). *Let $\rho \in \mathcal{P}$, and let $\delta > 0$. Let θ_δ be a smooth cut-off function such that $\frac{|\nabla\theta_\delta|^2}{\theta}$ is continuous and such that*

$$\begin{cases} \theta_\delta = 1 & \text{on } A_\delta := \bigcup_{i=1}^M B(X_i, \delta) \\ \theta_\delta = 0 & \text{outside } \bigcup_{i=1}^M B(X_i, 2\delta) \\ 0 \leq \theta_\delta \leq 1 & \text{on } \mathbb{R}^3 \\ \left| \frac{|\nabla\theta_\delta|^2}{\theta_\delta} - 2\Delta\theta_\delta \right| \leq K. \end{cases} \tag{4.2}$$

Let v be smooth and compactly supported function such that:

$$\begin{cases} \varepsilon\mathcal{T}(v) \leq K \\ \theta_\delta\rho + v \in \mathcal{P} \\ \text{dist}(\text{spt } v, \{X_1, \dots, X_M\}) \geq 3\delta. \end{cases} \tag{4.3}$$

Then, setting $Z = \sum_{1 \leq k \leq M} Z_k$, we have

$$G_\varepsilon^0(\theta_\delta\rho + v) \leq G_\varepsilon^0(\rho) + \varepsilon \left((1 + \varepsilon)K + \frac{Z}{\delta} \right).$$

Proof. We compute

$$\frac{|\nabla(\theta_\delta\rho)|^2}{\theta_\delta\rho} = \frac{|\theta_\delta\nabla\rho + \rho\nabla\theta_\delta|^2}{\theta_\delta\rho} = \theta_\delta \frac{|\nabla\rho|^2}{\rho} + \rho \frac{|\nabla\theta_\delta|^2}{\theta_\delta} + 2\nabla\rho \cdot \nabla\theta_\delta.$$

Concerning the second and third terms we remark that

$$\left| \int \rho \frac{|\nabla\theta_\delta|^2}{\theta_\delta} + 2\nabla\rho \cdot \nabla\theta_\delta \, dx \right| = \left| \int \left(\frac{|\nabla\theta_\delta|^2}{\theta_\delta} - 2\Delta\theta_\delta \right) \rho \, dx \right| \leq K.$$

Summarizing, we have obtained

$$\int \frac{|\nabla(\theta_\delta\rho)|^2}{4\theta_\delta\rho} \, dx \leq T(\rho) + K.$$

Since the support of v is away from that of $\theta_\delta\rho$ we have that

$$\varepsilon^2 T(\theta_\delta\rho + v) = \varepsilon^2 \mathcal{T}(\theta_\delta\rho) + \varepsilon^2 \mathcal{T}(v) \leq \varepsilon^2 T(\rho) + (\varepsilon^2 + \varepsilon)K.$$

Similarly,

$$\int V(x)\theta_\delta\rho \, dx \geq \int V(x)\rho \, dx - \int_{A_\delta^c} V(x)\rho \, dx \geq \int V(x)\rho \, dx - \frac{Z}{\delta},$$

and

$$\int V(x)v \, dx \geq 0.$$

Summing up the last inequalities gives the desired estimate. □

Example 4.2. We give for completeness an example of functions θ_δ and v satisfying the assumptions of Lemma 4.1 above. We define the real functions

$$f(t) = \begin{cases} e^{-1/t} & \text{if } t > 0 \\ 0 & \text{otherwise} \end{cases} \quad \text{and} \quad g(t) := \frac{f(t)}{f(t) + f(1-t)}.$$

When $\delta \ll \|X_i - X_j\|$ for $i \neq j$ one may consider

$$\theta_\delta(x) = \sum_{i=1}^M g \left(2 - \left(\frac{\|x - X_i\|}{\delta} \right)^2 \right).$$

Concerning v it is enough to consider any positive function h with smooth \sqrt{h} supported away from the points X_i and set $v = \beta h$ with β small enough so that $\theta_\delta \rho + v \in \mathcal{P}$: in that case the constant K in (4.3) does not depend on ε .

Lemma 4.3. *We have*

$$G^{0+}(\delta_{X_k}) \leq -\frac{Z_k^2}{4}.$$

Proof. It is enough to prove the inequality for X_1 and Z_1 and without loss of generality we may assume $X_1 = 0$. Consider a generic probability η and define

$$\rho_\varepsilon(x) = \eta^{1/\varepsilon}(x) = \frac{1}{\varepsilon^3} \eta(x/\varepsilon).$$

We have that ρ_ε weakly* converges to δ_0 , so that

$$G^{0+}(\delta_0) \leq \limsup_{\varepsilon \rightarrow 0} G_\varepsilon^0(\rho_\varepsilon) = T(\eta) - \int \frac{Z_1 \eta}{|x|} \, dx.$$

Taking the infimum with respect to η and using (4.1) gives what required. □

Theorem 4.4. *The limit functional G^0 exists and is given by the formula*

$$G^0(\rho) = -\frac{1}{4} \sum_{1 \leq k \leq M} Z_k^2 \rho(\{X_k\}). \tag{4.4}$$

Proof. We prove that

$$-\frac{1}{4} \sum_{1 \leq k \leq M} Z_k^2 \rho(\{X_k\}) \leq G^{0-}(\rho) \leq G^{0+}(\rho) \leq -\frac{1}{4} \sum_{1 \leq k \leq M} Z_k^2 \rho(\{X_k\}). \quad (4.5)$$

The last inequality in (4.5) follows by the convexity of the functional G^{0+} and Lemma 4.3. Indeed, we can write for every probability ρ

$$\rho = \sum_{1 \leq k \leq M} \alpha_k \delta_{X_k} + \alpha_0 \frac{\rho^\perp}{\alpha_0}$$

where

$$\alpha_k = \rho(\{X_k\}), \quad \rho^\perp = \rho \llcorner \{\mathbb{R}^3 \setminus \cup_{1 \leq k \leq M} \{X_k\}\}, \quad \alpha_0 = \rho^\perp(\mathbb{R}^3).$$

Since $\alpha_0 + \sum_{1 \leq k \leq M} \alpha_k = 1$ the convexity of G^+ gives

$$G^{0+}(\rho) \leq \sum_{1 \leq k \leq M} \alpha_k G^{0+}(\delta_{X_k}) + \alpha_0 G^{0+}(\rho^\perp / \alpha_0).$$

By Lemma 4.3 we have

$$G^{0+}(\delta_{X_k}) \leq -\frac{1}{4} Z_k^2$$

and, since $G^{0+}(\rho) = 0$ whenever ρ does not charge any of the points X_k (see Proposition 3.7), we have $G^{0+}(\rho^\perp / \alpha_0) = 0$ so that the desired inequality follows.

In order to prove the first inequality in (4.5) we have to show that for every ρ_ε weakly* converging to ρ we have

$$-\frac{1}{4} \sum_{1 \leq k \leq M} Z_k^2 \rho(\{X_k\}) \leq \liminf_{\varepsilon \rightarrow 0} G_\varepsilon^0(\rho_\varepsilon).$$

We apply Lemma 4.1 to ρ_ε with δ small enough and fixed, so that we can replace ρ_ε by

$$\theta_\delta \rho_\varepsilon + \nu_\varepsilon = \sum_{k=1}^M \theta_\delta \rho_\varepsilon \llcorner B(X_k, 2\delta) + \nu_\varepsilon = \sum_{k=1}^M \rho_\varepsilon^k + \nu_\varepsilon$$

where we defined $\rho_\varepsilon^k := \theta_\delta \rho_\varepsilon \llcorner B(X_k, 2\delta)$ and ν_ε is chosen as in Example 4.2 with a fixed function h so that the constant K does not depend on ε . We then have

$$\liminf_{\varepsilon \rightarrow 0} G_\varepsilon^0(\rho_\varepsilon) \geq \liminf_{\varepsilon \rightarrow 0} \left(G_\varepsilon^0 \left(\sum_{k=1}^M \rho_\varepsilon^k + \nu_\varepsilon \right) - \varepsilon K_\delta \right) = \liminf_{\varepsilon \rightarrow 0} G_\varepsilon^0 \left(\sum_{k=1}^M \rho_\varepsilon^k + \nu_\varepsilon \right),$$

where we can take $K_\delta = 2K + \frac{Z}{\delta}$ for $\varepsilon \leq 1$ and

$$G_\varepsilon^0 \left(\sum_{k=1}^M \rho_\varepsilon^k + \nu_\varepsilon \right) = \sum_{k=1}^M \left(\varepsilon^2 T(\rho_\varepsilon^k) - \varepsilon Z_k U_{X_k}(\rho_\varepsilon^k) \right) + \varepsilon^2 T(\nu_\varepsilon) - \varepsilon U(\nu_\varepsilon).$$

Concerning the first M terms we infer from (4.1) that

$$\begin{aligned} \varepsilon^2 T(\rho_\varepsilon^k) - \varepsilon Z_k U_{X_k}(\rho_\varepsilon^k) &= \left(\varepsilon^2 T \left(\frac{\rho_\varepsilon^k}{\int d\rho_\varepsilon^k} \right) - \varepsilon Z_k U_{X_k} \left(\frac{\rho_\varepsilon^k}{\int d\rho_\varepsilon^k} \right) \right) \int d\rho_\varepsilon^k \\ &\geq -\frac{Z_k^2}{4} \int d\rho_\varepsilon^k. \end{aligned}$$

As $\varepsilon \rightarrow 0$ we have

$$\lim_{\varepsilon \rightarrow 0} \int d\rho_\varepsilon^k = \int_{B(X_k, 2\delta)} \theta_\delta \rho \leq \rho(B(X_k, 2\delta)).$$

The last term $\varepsilon^2 T(\nu_\varepsilon) - \varepsilon U(\nu_\varepsilon)$ vanishes as $\varepsilon \rightarrow 0$. Summing up we obtained

$$\liminf_{\varepsilon \rightarrow 0} G_\varepsilon^0 \left(\sum_{k=1}^M \rho_\varepsilon^k + \nu_\varepsilon \right) \geq -\sum_{k=1}^M \frac{Z_k^2}{4} \rho(B(X_k, 2\delta)).$$

Letting now $\delta \rightarrow 0$ gives the desired inequality. □

Remark 4.5. Since the correlation term C is non-negative, we obtain from the preceding that

$$\forall \rho \in \mathcal{P}, \quad G^-(\rho) \geq G^{0-}(\rho) = -\frac{1}{4} \sum_{1 \leq k \leq M} Z_k^2 \rho(\{X_k\}).$$

which is a more precise lower estimate of G^- than that obtained in Proposition 3.7.

Remark 4.6. The explicit form of G^0 given in (4.4) allows to directly deduce that

$$\operatorname{argmin} G^0 = \left\{ \rho = \sum_{i \in I_{\max}} \alpha_i \delta_{X_i} : 0 \leq \alpha_i \leq 1, \sum \alpha_i = 1 \right\},$$

where I_{\max} denotes the set of indices i such that $Z_i = \max\{Z_1, \dots, Z_M\}$. In other words, the optimal configurations for the limit functional G^0 concentrate on the points X_i having the highest nuclei charges Z_i . We shall see in Remark 5.10 below that the situation is somewhat more intricate in the interacting case $b > 0$.

5. The interacting case for $N = 2$

The complete characterization of the functional G defined as the Γ -limit of the functionals G_ε as $\varepsilon \rightarrow 0$, in the general case $M \geq 1$ and $N \geq 1$ seems a very difficult issue, mainly due to the general form of the localization Lemma 5.1 below, that is at the moment unavailable. We then limit ourselves in this section to consider the case $N = 2$ that we can handle completely.

For every $\rho \in \mathcal{P}^-$ we denote by $\rho^\#$ the atomic measure

$$\rho^\# = \sum_{i=1}^M \rho(\{X_i\})\delta_{X_i},$$

where X_i are the (fixed) positions of the nuclei ($i = 1, \dots, M$). We also denote by ρ^\perp the measure

$$\rho^\perp = \rho - \rho^\#$$

that does not charge any of the points X_i . In other words, $\rho^\#$ and ρ^\perp are respectively the restrictions of ρ to the sets $\{X_1, \dots, X_M\}$ and to its complement.

The following Lemma extends the localization argument of Lemma 4.1 in presence of the correlation term C .

Lemma 5.1 (Localization). *Let $N = 2$, $\rho \in \mathcal{P}$, and $\delta > 0$, and let θ_δ and v be as in Lemma 4.1. Assume further that $v = v^1 + v^2$ with $\|v^1\| = \|v^2\|$ and $\text{spt}(v^1) \cap \text{spt}(v^2) = \emptyset$. Then we have*

$$G_\varepsilon(\theta_\delta \rho + v) \leq G_\varepsilon(\rho) + \varepsilon \left((1 + \varepsilon)K + \frac{Z + 2b}{\delta} + \frac{b}{\text{dist}(\text{spt}(v^1), \text{spt}(v^2))} \right).$$

Proof. By Lemma 4.1 we only need the following estimate of the transport term C :

$$C(\theta_\delta \rho + v) \leq C(\rho) + \frac{2}{\delta} + \frac{1}{\text{dist}(\text{spt}(v^1), \text{spt}(v^2))}.$$

It is convenient to introduce the set $R = \mathbb{R}^3 \setminus A_\delta$, being A_δ defined in (4.2).

We denote by P an optimal transport plan for ρ , which is also symmetric with respect to a permutation of the variables and we define a new transport plan \tilde{P} as below.

$$\begin{aligned} \tilde{P}^1 &= \tilde{P}|_{A_\delta \times A_\delta} = \min\{\theta_\delta(x), \theta_\delta(y)\} P|_{A_\delta \times A_\delta}, \\ \tilde{P}^2 &= \tilde{P}|_{A_\delta \times R} = (\theta_\delta \rho - \pi_1^\# \tilde{P}^1) \otimes \frac{v}{\|v\|}, \\ \tilde{P}^3 &= \tilde{P}|_{R \times A_\delta} = \frac{v}{\|v\|} \otimes (\theta_\delta \rho_\varepsilon - \pi_2^\# \tilde{P}^1), \\ \tilde{P}^4 &= 2 \left(1 - \frac{\|\theta_\delta \rho_\varepsilon - \pi_2^\# \tilde{P}^1\|}{\|v\|} \right) (v^1 \otimes v^2 + v^2 \otimes v^1). \end{aligned}$$

The fact that $\tilde{P}^2 \geq 0$ (and similarly that $\tilde{P}^3 \geq 0$) follows from $\tilde{P}^1 \leq \theta_\delta(x)P|_{A_\delta \times A_\delta}$. To check that $\tilde{P}^4 \geq 0$ it is enough to compute

$$\begin{aligned} \|v\| - \|\theta_\delta \rho - \pi_2^\# \tilde{P}^1\| &= 1 - \|\theta_\delta \rho\| - (\|\theta_\delta \rho\| - \|\pi_2^\# \tilde{P}^1\|) \\ &= 1 + \|\pi_2^\# \tilde{P}^1\| - \|\theta_\delta \rho\| - \|\theta_\delta \rho\| \\ &= \int [1 + \min\{\theta_\delta(x), \theta_\delta(y)\} - \theta_\delta(x) - \theta_\delta(y)] dP(x, y) \geq 0 \end{aligned}$$

where the last inequality follows since the integrand is non-negative. Also note that

$$v - \pi_1^\# \tilde{P}^3 = \left(1 - \frac{\|\theta_\delta \rho_\varepsilon - \pi_2^\# \tilde{P}^1\|}{\|v\|}\right) v$$

so that $\pi_1^\# \tilde{P}^4 = v - \pi_1^\# \tilde{P}^3$.

In order to show that the two marginals of \tilde{P} coincide with $\theta_\delta \rho + v$, since \tilde{P} is symmetric, it is enough to check the first marginal. Since R is the complement of A_δ we compute the restriction of the marginal to these two sets. Then

$$\begin{aligned} \text{on } A_\delta \text{ we have } \pi_1^\# \tilde{P} &= \pi_1^\# \tilde{P}^1 + \pi_1^\# \tilde{P}^2 = \pi_1^\# \tilde{P}^1 + \theta_\delta \rho - \pi_1^\# \tilde{P}^1 = \theta_\delta \rho, \\ \text{on } R \text{ we have } \pi_1^\# \tilde{P} &= \pi_1^\# \tilde{P}^3 + \pi_1^\# \tilde{P}^4 = \pi_1^\# \tilde{P}^3 + v - \pi_1^\# \tilde{P}^3 = v. \end{aligned}$$

Since the quantity $\int_{(\mathbb{R}^3)^2} c dP$ to be minimized in C is linear with respect to P it is enough to estimate it for each of the components of \tilde{P} above and using the facts that

$$\begin{cases} \text{dist}(\text{spt}(v^1), \text{spt}(v^2)) > 0 \\ 0 \leq \min\{\theta_\delta(x), \theta_\delta(y)\} \leq 1 \\ \text{dist}(\text{spt } v, \{X_1, \dots, X_M\}) \geq 3\delta, \end{cases}$$

we obtain

$$\int_{(\mathbb{R}^3)^2} c(x, y) d\tilde{P}(x, y) \leq \int_{(\mathbb{R}^3)^2} c(x, y) dP(x, y) + \frac{1}{\delta} + \frac{1}{\delta} + \frac{1}{\text{dist}(\text{spt}(v^1), \text{spt}(v^2))}$$

which is the desired inequality. □

We are now in position to prove both the existence of the Γ -limit of the functionals G_ε and a property of it that will be very useful in the following to obtain an explicit representation formula.

Theorem 5.2. *For every $\rho \in \mathcal{P}^-$ the Γ -limit G of the functionals G_ε exists and we have*

$$G(\rho) = G(\rho^\#).$$

Proof. By the compactness of the Γ -convergence, the Γ -limit G exists, at least for a subsequence $\varepsilon_n \rightarrow 0$; as stated in Section 3, since later we will characterize this Γ -limit explicitly, we may assume it does not depend on the subsequence, so that the entire family G_ε Γ converges and $G^- = G^+ = G$. Let $\rho \in \mathcal{P}^-$. Writing $\rho = \rho^\# + \rho^\perp$, it is then enough to show the inequalities below:

$$\begin{aligned} G^-(\rho^\# + \rho^\perp) &\leq G^+(\rho^\#), \\ G^+(\rho^\# + \rho^\perp) &\geq G^-(\rho^\#). \end{aligned} \tag{5.1}$$

Let us prove the first inequality in (5.1) for $\rho \in \mathcal{P}$ in the special case that $\rho^\perp = \rho_1^\perp + \rho_2^\perp$ with $\|\rho_1^\perp\| = \|\rho_2^\perp\|$, ρ_i^\perp smooth, with disjoint and compact supports; in addition we assume that $\text{dist}(\text{spt } \rho^\#, \text{spt } \rho^\perp) > 0$. Denote by ρ_ε a family weakly* converging to $\rho^\#$ and such that

$$G^+(\rho^\#) = \limsup_{\varepsilon \rightarrow 0} G_\varepsilon(\rho_\varepsilon).$$

Define, for δ small enough and θ_δ as in Lemma 4.1, $\tilde{\rho}_\varepsilon = \theta_\delta \rho_\varepsilon + v_\varepsilon$, where we choose $v_\varepsilon = a_\varepsilon \rho^\perp = a_\varepsilon \rho_1^\perp + a_\varepsilon \rho_2^\perp$ and a_ε is such $\tilde{\rho}_\varepsilon \in \mathcal{P}$. Then the assumptions of Lemma 5.1 are satisfied. Then $\tilde{\rho}_\varepsilon \xrightarrow{*} \theta_\delta \rho^\# + \rho^\perp$ and we have by Lemma 5.1

$$G_\varepsilon(\tilde{\rho}_\varepsilon) \leq G_\varepsilon(\rho_\varepsilon) + \varepsilon K_\delta,$$

where the constant K_δ only depends on δ . Passing to the limit as $\varepsilon \rightarrow 0$ we have

$$G^-(\theta_\delta \rho^\# + \rho^\perp) \leq G^+(\rho^\#).$$

Using now the lower semicontinuity of G^- we have, as $\delta \rightarrow 0$

$$G^-(\rho^\# + \rho^\perp) \leq G^+(\rho^\#).$$

In order to extend the inequality above to the general case of $\rho \in \mathcal{P}^-$ with general ρ^\perp we use again the lower semicontinuity property of G^- and a density argument.

To prove the second inequality in (5.1) we argue in a similar way: take as ρ_ε a family weakly* converging to $\rho = \rho^\# + \rho^\perp$ and such that

$$G^+(\rho^\# + \rho^\perp) = \limsup_{\varepsilon \rightarrow 0} G_\varepsilon(\rho_\varepsilon).$$

Thanks to Lemma 5.1, we may construct $\tilde{\rho}_\varepsilon = \theta_\delta \rho_\varepsilon + v_\varepsilon$ as above, with the cut-off function θ_δ as in (4.2) and taking, for example,

$$v_\varepsilon(x) = a_\varepsilon \left[h \left(x - \frac{x_0}{\varepsilon} \right) + h \left(x + \frac{x_0}{\varepsilon} \right) \right],$$

with $h \geq 0$ smooth and compactly supported, $x_0 \neq 0$ and a_ε suitably chosen so that $\tilde{\rho}_\varepsilon \in \mathcal{P}$. We get $v_\varepsilon \xrightarrow{*} 0$ and

$$G_\varepsilon(\tilde{\rho}_\varepsilon) \leq G_\varepsilon(\rho_\varepsilon) + \varepsilon K_\delta.$$

Then, passing to the limit as $\varepsilon \rightarrow 0$, we have

$$G^+(\rho^\# + \rho^\perp) \geq G^-(\theta_\delta \rho);$$

passing now to the limit as $\delta \rightarrow 0$ gives $G^+(\rho^\# + \rho^\perp) \geq G^-(\rho^\#)$ as required. \square

By Theorem 5.2 all Γ -limits of subsequences G_{ε_n} depend only on $\rho^\#$. In Theorem 5.7 we will characterize by an explicit formula the Γ -limit of G_{ε_n} independently of the subsequence ε_n , obtaining in this way the Γ -limit of the whole family G_ε .

The following definition of partial or fractional transport cost appeared in the w^* relaxation \overline{C} of C in equation (2.3) and will appear in the formula for the Γ -limit:

$$C(\rho, 2\alpha - 1) = \begin{cases} \min \{ \mathcal{C}(\mu) : 0 \leq \mu \leq \rho, \|\mu\| = 2\alpha - 1 \} & \text{if } \alpha \geq 1/2 \\ 0 & \text{otherwise.} \end{cases}$$

Definition 5.3. For all $b, Z > 0$ and $0 \leq \alpha \leq 1$ we define

$$g_b(Z, \alpha) := \inf_{\|\rho\|=\alpha} \{ \mathcal{T}(\rho) + bC(\rho, 2\alpha - 1) - Z\mathcal{U}_0(\rho) \}. \tag{5.2}$$

Remark 5.4. Using again the different homogeneities of the three addenda with respect to the rescaling of measures we have

$$g_b(Z, \alpha) = -\frac{1}{4} \sup_{\|\rho\|=\alpha} \frac{(Z\mathcal{U}_0(\rho) - bC(\rho, 2\alpha - 1))_+^2}{\mathcal{T}(\rho)}$$

so that for $\alpha \leq 1/2$ the equality

$$g_b(Z, \alpha) = -\frac{1}{4}\alpha Z^2$$

holds. Moreover from (5.2), again rescaling the measures, for every $\varepsilon > 0$ we have

$$\varepsilon^2 \mathcal{T}(\rho) + \varepsilon bC(\rho, 2\alpha - 1) - \varepsilon Z\mathcal{U}_0(\rho) \geq g_b(Z, \alpha)$$

for all $\rho \in \mathcal{P}^-$ with $\|\rho\| = \alpha$.

It is clear from the definition of g_b that it is concave non-increasing in Z , and we shall prove in Lemma 6.3 (using the other equivalent Definition 6.1) that it is convex non-increasing in α . Discussing the existence of minimizers is out of the scope of this paper, however we will need some almost optimal measures which we study in the next proposition.

Proposition 5.5. For all $\lambda > 0$ there exist $r = r(\lambda)$ and $\rho \in \mathcal{P}^-$ with $\|\rho\| = \alpha$ such that $\text{spt}(\rho) \subset B(0, r)$ and

$$\mathcal{T}(\rho) + bC(\rho, 2\alpha - 1) - Z\mathcal{U}_0(\rho) \leq g_b(Z, \alpha) + \lambda.$$

Proof. This is an indirect variant of Lemma 5.1 in which the fractional transport cost appears and $\varepsilon = 1$. So we carefully apply Lemmas 4.1 and 5.1. We start from $\tilde{\rho} \in \mathcal{P}^-$ such that $\|\tilde{\rho}\| = \alpha$ and

$$\mathcal{T}(\tilde{\rho}) + bC(\tilde{\rho}, 2\alpha - 1) - Z\mathcal{U}_0(\tilde{\rho}) \leq g_b(Z, \alpha) + \frac{\lambda}{3}.$$

We need to modify $\tilde{\rho}$ so that the support becomes compact. Let also $\tilde{\mu} \leq \tilde{\rho}$ with $\|\tilde{\mu}\| = 2\alpha - 1$ and $\mathcal{C}(\tilde{\mu}) = C(\tilde{\rho}, 2\alpha - 1)$. Consider θ_δ and ν as in Lemma 5.1 with δ large enough so that we may have K small and

$$2K + \frac{Z + 2b}{\delta} + \frac{b}{\text{dist}(\text{spt}(\nu_\varepsilon^1), \text{spt}(\nu_\varepsilon^2))} \leq \frac{\lambda}{3}.$$

Here we set $\rho = \theta_\delta \tilde{\rho} + \nu$ and assume $\|\rho\| = \alpha$ (instead of $\rho \in \mathcal{P}$ as in Lemma 5.1). Then

$$\int \nu \, dx = \alpha - \|\theta_\delta \tilde{\rho}\|.$$

By Lemma 4.1 with $\varepsilon = 1$ we have

$$\mathcal{T}(\rho) - Z\mathcal{U}_0(\rho) \leq \mathcal{T}(\tilde{\rho}) - Z\mathcal{U}_0(\tilde{\rho}) + \frac{\lambda}{3}.$$

To estimate the fractional transport term $C(\rho, 2\alpha - 1)$ we consider $\mu = \theta_\delta \tilde{\mu} + \beta_0 \nu$ where β_0 is such that the total variation of μ is equal to $2\alpha - 1$. To show that $\mu \leq \rho$ we need to show that $\beta_0 \leq 1$. This is equivalent to say that

$$2\alpha - 1 - \int \theta_\delta d\tilde{\mu} \leq \int d\nu = \alpha - \int \theta_\delta d\tilde{\rho}$$

which is the inequality

$$\int \theta_\delta d(\tilde{\rho} - \tilde{\mu}) \leq 1 - \alpha = \int d(\tilde{\rho} - \tilde{\mu}).$$

We then apply the transport estimate of Lemma 5.1, up to a rescale of the measures, to get

$$C(\rho, 2\alpha - 1) \leq \mathcal{C}(\mu) \leq \mathcal{C}(\tilde{\mu}) + \frac{\lambda}{3} = C(\tilde{\rho}, 2\alpha - 1) + \frac{\lambda}{3}.$$

which concludes the proof. □

A short investigation of the structure of certain optimal transport plans will be used in the next theorem.

Lemma 5.6. *Let $X_0, \dots, X_M \in \mathbb{R}^d$ and let $\delta > 0$ be such that $\delta \ll \min_{i,j} |X_i - X_j|$. Consider $\rho = \sum_{i=0}^M \alpha_i \rho_i$ with ρ_i probability measures such that $\text{spt } \rho_i \subset B(X_i, \delta)$ and $\sum_{i=0}^M \alpha_i = 1$. Let P be an optimal transport plan for ρ , then*

$$\begin{cases} P(B(X_i, \delta) \times B(X_i, \delta)) = 0 & \text{if } \alpha_i \leq 1/2 \\ P(B(X_i, \delta) \times B(X_i, \delta)) = 2\alpha_i - 1 & \text{if } \alpha_i > 1/2. \end{cases}$$

It follows that there exists K depending only on $\min_{i \neq j} |X_i - X_j| - 2\delta$ such that the following alternative holds:

- (1) $C(\rho) \leq K$ if $\alpha_i \leq 1/2$ for all i ;
- (2) If $\alpha_i > 1/2$ for some i then

$$C(\alpha_i \rho_i, 2\alpha_i - 1) \leq C(\rho) \leq C(\alpha_i \rho_i, 2\alpha_i - 1) + K.$$

Note that in the above statement we add a point X_0 , not corresponding to a nucleus, to the points X_1, \dots, X_M : this will be handy in the proof of Theorem 5.7 below.

Proof. Let $i \in \{0, \dots, M\}$ and let $l_1 := P(B(X_i, \delta) \times B(X_i, \delta))$, we may compute

$$\begin{aligned} P((B(X_i, c\delta) \times \mathbb{R}^d) \cup (\mathbb{R}^d \times B(X_i, \delta))) &= P(B(X_i, \delta) \times \mathbb{R}^d) + P(\mathbb{R}^d \times B(X_i, \delta)) \\ &\quad - P(B(X_i, \delta) \times B(X_i, \delta)) \leq 2\alpha_i - l_1. \end{aligned}$$

For the case $\alpha_i \leq 1/2$, if $l_1 > 0$ then, since $2\alpha_i - l_1 < 1$, there exist $j, k \neq i$ such that

$$l_2 := P(B(X_j, \delta) \times B(X_k, \delta)) > 0.$$

Define $s = \min\{l_1, l_2\}$ and

$$P_1 = \frac{s}{l_1} P_{|B(X_i, \delta) \times B(X_i, \delta)}, \quad P_2 = \frac{s}{l_2} P_{|B(X_j, \delta) \times B(X_k, \delta)},$$

and rewrite $P = P_1 + P_2 + P_R$ where P_R is defined by this same equality. Since the quantity $c(P) = \int_{(\mathbb{R}^3)^2} c \, dP$ is linear in P we have

$$c(P) = c(P_1) + c(P_2) + c(P_R) \geq \frac{s}{2\delta} + c(P_R).$$

Define the possibly non-symmetric competitor \tilde{P} by

$$\tilde{P} = \frac{1}{s} \left(\pi_1^\# P_1 \otimes \pi_2^\# P_2 + \pi_1^\# P_2 \otimes \pi_2^\# P_1 \right) + P_R,$$

so that it has the same marginal as P . Concerning the transportation cost we have

$$c(\tilde{P}) \leq \frac{s}{|X_i - X_k| - 2\delta} + \frac{s}{|X_j - X_i| - 2\delta} + c(P_R)$$

which is smaller than $c(P)$ for δ as in the assumptions, and this contradicts the optimality of P .

Analogously, if $\alpha_i > 1/2$, since $\rho(\mathbb{R}^d \setminus B(X_i, \delta)) = 1 - \alpha_i$ we have

$$\begin{aligned} l_1 &= P(B(X_i, \delta) \times B(X_i, \delta)) \\ &= P(B(X_i, \delta) \times \mathbb{R}^d) - P(B(X_i, \delta) \times (\mathbb{R}^d \setminus B(X_i, \delta))) \\ &\geq \alpha_i - (1 - \alpha_i) = 2\alpha_i - 1. \end{aligned}$$

The strict inequality would imply again that $2\alpha_i - l_1 < 1$ and then again there exist $j, k \neq i$ such that $P(B(X_j, \delta) \times B(X_k, \delta)) > 0$. This would contradict the optimality of P as in the first case.

To deduce (5.6) we consider an optimal transport plan P_{opt} and remark that for x in the support of P_{opt} for all i, j

$$\frac{1}{|x_i - x_j|} \leq \frac{1}{\min_{i \neq j} |X_i - X_j| - 2\delta}$$

so that we can take

$$K = \frac{N(N - 1)}{2 \min_{i \neq j} |X_i - X_j| - 2\delta}.$$

To prove the first inequality in (5.6) consider a symmetric optimal transport plan P for $C(\rho)$. Let $P' = P \lfloor B(X_i, \delta) \times B(X_i, \delta)$ and let $\mu := \pi_1^\# P' (= \pi_2^\# P')$. Clearly $0 \leq \mu \leq \alpha_i \rho_i$ and $\|\mu\| = \|P'\| = 2\alpha_i - 1$, then

$$C(\alpha_i \rho_i, 2\alpha_i - 1) \leq c(P') \leq c(P') + c(P - P') = C(\rho).$$

For the second inequality in (5.6), let $\mu \leq \alpha_i \rho_i$ with $\|\mu\| = 2\alpha_i - 1$ be such that

$$C(\mu) = C(\alpha_i \rho_i, 2\alpha_i - 1),$$

and let P_μ be an optimal plan for μ . Consider an optimal plan P_R for $\rho - \mu$. The plan $P = P_\mu + P_R \in \Pi(\rho)$ and then

$$C(\rho) \leq c(P) = c(P_\mu) + c(P_R) = C(\mu) + C(\rho - \mu) = C(\alpha_i \rho_i, 2\alpha_i - 1) + C(\rho - \mu).$$

We conclude observing that

$$C(\rho - \mu) = \|\rho - \mu\| C\left(\frac{\rho - \mu}{\|\rho - \mu\|}\right) \leq K \|\rho - \mu\| \leq K$$

since

$$\rho - \mu = \alpha_0 \rho_0 + \dots + (1 - \alpha_i) \rho_i + \dots + \alpha_M \rho_M$$

and

$$\alpha_0, \dots, 1 - \alpha_i, \dots, \alpha_M \leq \frac{1}{2} < \alpha_i = \frac{\|\rho - \mu\|}{2}$$

so (5.6) applies to $\frac{\rho - \mu}{\|\rho - \mu\|}$. □

By Theorem 5.2 we may now focus on the formula for $G(\rho^\#)$.

Theorem 5.7. *Let $\rho \in \mathcal{P}^-$ be such that*

$$\rho^\# = \sum_{i=1}^M \alpha_i \delta_{X_i} \quad \text{with } \alpha_i \geq 0 \text{ and } \sum_{i=1}^M \alpha_i \leq 1.$$

Then the following formula holds:

$$G(\rho) = \sum_{i=1}^M g_b(Z_i, \alpha_i).$$

Proof. It follows from Theorem 5.2 that $G(\rho) = G(\rho^\#)$, so we shall prove the result for $\rho = \rho^\#$.

Let $\delta > 0$ be such that $\delta \ll \min_{i,j} |X_i - X_j|$. We start with the $\Gamma - \lim \sup$ inequality. For every $\varepsilon > 0$ we add to the M -uple (X_1, \dots, X_M) an additional point Y_ε such that

$$\lim_{\varepsilon \rightarrow 0} \|Y_\varepsilon\| \rightarrow \infty.$$

Let $\lambda > 0$, let ρ_i be a measure with compact support obtained from Proposition 5.5 applied with parameters $\lambda_i = \frac{\lambda}{M}, \alpha_i, Z_i$. Let $h \in C_0^\infty(B(0, \delta))$ a positive function such that

$$\int h \, dx = 1 \quad \text{and} \quad \int \frac{|\nabla h|^2}{h} \, dx < +\infty$$

and define

$$\rho_\varepsilon^0(x) = \alpha_0 h(x - Y_\varepsilon) \quad \text{with } \alpha_0 = 1 - \sum_{i=1}^M \alpha_i.$$

For $i = 1, \dots, M$ let

$$\rho_\varepsilon^i(x) = \rho_i^{1/\varepsilon}(x - X_i) = \frac{1}{\varepsilon^3} \rho_i \left(\frac{x - X_i}{\varepsilon} \right) \quad \text{and} \quad \rho_\varepsilon(x) = \sum_{i=0}^M \rho_\varepsilon^i(x).$$

For ε small enough the supports of ρ_ε^0 and ρ_ε^i are contained in $B(Y_\varepsilon, \delta)$ and $B(X_i, \delta)$ respectively. We estimate $G_\varepsilon(\rho_\varepsilon)$ from above. Since

$$T(\rho_\varepsilon) = \sum_{i=1}^M T(\rho_\varepsilon^i) + \alpha_0 T(h),$$

$$U(\rho_\varepsilon) \geq \sum_{i=1}^M Z_i \mathcal{U}_{X_i}(\rho_\varepsilon^i),$$

we deduce

$$G_\varepsilon(\rho_\varepsilon) \leq \sum_{i=1}^M (\varepsilon^2 T(\rho_\varepsilon^i) - \varepsilon Z_i \mathcal{U}_{X_i}(\rho_\varepsilon^i)) + \varepsilon^2 \alpha_0 T(h) + \varepsilon C(\rho_\varepsilon). \tag{5.3}$$

We then need to decompose $C(\rho_\varepsilon)$. By Lemma 5.6 if $\alpha_i \leq 1/2$ for all i then $C(\rho_\varepsilon) \leq K$ and passing to the lim sup

$$G^+(\rho^\#) \leq \limsup_{\varepsilon \rightarrow 0} G_\varepsilon(\rho_\varepsilon) \leq \sum_i g_b(Z_i, \alpha_i) + \lambda$$

because the last two terms in (5.3) go to 0, and the first term is computed by the homogeneity of the energy and the choice of ρ_i . If for one $i \in \{1, \dots, M\}$, $\alpha_i > 1/2$, we assume without loss of generality that it is α_1 , then by Lemma 5.6

$$C(\rho_\varepsilon) \leq C(\rho_\varepsilon^1, 2\alpha_1 - 1) + K.$$

Then

$$\begin{aligned} G_\varepsilon(\rho_\varepsilon) &\leq \varepsilon^2 T(\rho_\varepsilon^1) - \varepsilon U_{Z_1}(\rho_\varepsilon^1) + \varepsilon C_2(\rho_\varepsilon^1, 2\alpha_1 - 1) \\ &\quad + \sum_{i=2}^M (\varepsilon^2 T(\rho_\varepsilon^i) - \varepsilon U_{Z_i}(\rho_\varepsilon^i)) + \varepsilon^2 \alpha_0 T(h) + \varepsilon K \end{aligned}$$

and again we conclude by the homogeneity of the energy and the choice of ρ_i . The case $\alpha_0 > 1/2$ can be excluded by considering a second sequence $\tilde{Y}_\varepsilon = -Y_\varepsilon$ and then defining

$$\rho_\varepsilon^0 = \frac{\alpha_0}{2} h(x - Y_\varepsilon) + \frac{\alpha_0}{2} h(x - \tilde{Y}_\varepsilon),$$

which has the same properties needed in the proof but do not concentrate too much mass in a ball of radius δ : we are then applying Lemma 5.6 with the $M + 2$ points $X_1, \dots, X_M, Y_\varepsilon, \tilde{Y}_\varepsilon$.

We now deal with the $\Gamma - \lim \inf$ inequality. Let $\rho_\varepsilon \xrightarrow{*} \rho^\#$. By Lemma 5.1 we can replace ρ_ε by $\theta_\delta \rho_\varepsilon + \nu_\varepsilon$ with ν_ε chosen as in Example 4.2 so that the constant K does not depend on δ , since

$$\liminf_{\varepsilon \rightarrow 0} G_\varepsilon(\theta_\delta \rho_\varepsilon + \nu_\varepsilon) \leq \liminf_{\varepsilon \rightarrow 0} G_\varepsilon(\rho_\varepsilon).$$

We denote by $\rho_\varepsilon^i := \theta_\delta \rho_\varepsilon|_{B(X_i, 2\delta)}$ and we have, for some constant K_δ that does not depend on ε , the inequality

$$G_\varepsilon(\rho_\varepsilon) \geq \sum_{i=1}^M (\varepsilon^2 T(\rho_\varepsilon^i) - \varepsilon U_{Z_i}(\rho_\varepsilon^i)) + \varepsilon^2 \alpha_0 T(\nu_\varepsilon) + \varepsilon C(\rho_\varepsilon) - \varepsilon \frac{K}{2\delta}.$$

Again we need to look at $C(\rho_\varepsilon)$. If $\alpha_i \leq 1/2$ for all i we just use that $C(\rho_\varepsilon) \geq 0$ and get

$$G_\varepsilon(\rho_\varepsilon) \geq -\frac{1}{4} \sum_{i=1}^M Z_i^2 \|\rho_\varepsilon^i\| + \varepsilon^2 \alpha_0 T(v_\varepsilon) - \varepsilon \frac{K}{2\delta}.$$

When $\varepsilon \rightarrow 0$ and then $\delta \rightarrow 0$ we obtain

$$\liminf_{\varepsilon \rightarrow 0} G_\varepsilon(\rho_\varepsilon) \geq -\frac{1}{4} \sum_{i=1}^M Z_i^2 \int_{B(X_i, 2\delta)} \rho dx \xrightarrow{\delta \rightarrow 0} -\frac{1}{4} \sum_{i=1}^M Z_i^2 \alpha_i = \sum_i g_b(Z_i, \alpha_i)$$

which concludes the proof in this case.

If $\alpha_1 > 1/2$ (or any α_i up to reindexing) then for ε small enough $\|\rho_\varepsilon^1\| = \rho_\varepsilon^1(B(X_1, 2\delta)) > 1/2$ and then by Lemma 5.6

$$G_\varepsilon(\rho_\varepsilon) \geq \varepsilon^2 T(\rho_\varepsilon^1) + \varepsilon C(\rho_\varepsilon^1, 2\|\rho_\varepsilon^1\| - 1) - \varepsilon U_{Z_1}(\rho_\varepsilon^1) + \sum_{i=2}^M (\varepsilon^2 T(\rho_\varepsilon^i) - \varepsilon U_{Z_i}(\rho_\varepsilon^i)) + \varepsilon^2 \alpha_0 T(v_\varepsilon) + \varepsilon \frac{K}{2\delta}.$$

By the homogeneity of the three terms

$$G_\varepsilon(\rho_\varepsilon) \geq g_b(Z_1, \|\rho_\varepsilon^1\|) - \frac{1}{4} \sum_{i=2}^M Z_i^2 \|\rho_\varepsilon^i\| + \varepsilon^2 \alpha_0 T(v_\varepsilon) + \varepsilon \frac{K}{2\delta}.$$

Passing again to the limit for $\varepsilon \rightarrow 0$ and then $\delta \rightarrow 0$, and using the lower semi-continuity of $g_b(Z_i, \cdot)$ allows to conclude in this case. □

Remark 5.8. Since $\sum_i \alpha_i \leq 1$ at most one of the α_i may be greater than $1/2$ and we will always assume that is the first α_1 . Then, recalling that by definition,

$$\rho = \rho^\# + \rho^\perp = \sum_{i=1}^M \alpha_i \delta_{X_i} + \rho^\perp \quad \forall \rho \in \mathcal{P}^-,$$

we have two possible cases:

$$G(\rho) = G(\rho^\#) = -\frac{1}{4} \sum_{i=1}^M \alpha_i Z_i^2, \quad \text{if } \alpha_1 \leq \frac{1}{2},$$

or

$$G(\rho) = G(\rho^\#) = g_b(Z_1, \alpha_1) - \frac{1}{4} \sum_{i=2}^M \alpha_i Z_i^2, \quad \text{if } \alpha_1 > \frac{1}{2}.$$

We are now in a position to study the minimization problem

$$\min \{G(\rho) : \rho \in \mathcal{P}^-\}. \tag{5.4}$$

Theorem 5.9. *Let us assume that $M \geq 2$. Then the minimization problem (5.4) has a solution $\rho \in \mathcal{P}^-$. Moreover, every such a minimizer ρ belongs to \mathcal{P} and is of the form*

$$\rho = \rho^\# = \sum_{i=1}^M \alpha_i \delta_{X_i} \quad \text{with} \quad \sum_{i=1}^M \alpha_i = 1.$$

Note that the case $M = 1$ is discussed in Remark 6.6 below.

Proof. The existence of an optimal ρ follows from the weak* compactness of \mathcal{P}^- and lower semicontinuity of the Γ -limit G . For such a ρ , set $\alpha_i = \rho(\{X_i\})$. Then, by Theorem 5.7, $\rho^\# := \sum_{i=1}^M \alpha_i \delta_{X_i}$ is also optimal and we claim that $\sum_{i=1}^M \alpha_i = 1$. Indeed if $\sum_{i=1}^M \alpha_i < 1$, then there exists j such that $\alpha_j < 1/2$ and we may consider $\bar{\rho} := \rho + \eta \delta_{X_j}$ where η is such that

$$0 < \eta < \min \left\{ \frac{1}{2} - \alpha_j, 1 - \sum \alpha_i \right\}.$$

Then, by applying again Theorem 5.7, we obtain

$$\begin{aligned} G(\bar{\rho}) &= G(\rho^\# + \eta \delta_{X_j}) = -\frac{(\alpha_j + \eta)}{4} Z_j^2 + \sum_{i \neq j} g_b(Z_i, \alpha_i) \\ &= -\frac{1}{4} \eta Z_j^2 + G(\rho^\#) < G(\rho^\#) \end{aligned}$$

where we used the fact that $\alpha_j + \eta < 1/2$. □

Remark 5.10. To illustrate the previous analysis, we now discuss the structure of the minimizers of problem (5.4) in the special case $N = M = 2$. Without loss of generality we may assume that $Z_1 \geq Z_2$, then it follows from Theorem 5.9 that the minimizers of (5.4) are of the form

$$\rho = \alpha \delta_{X_1} + (1 - \alpha) \delta_{X_2}$$

with $\alpha \in [0, 1]$ minimizing the problem

$$\min \{ g_b(Z_1, \alpha) + g_b(Z_2, 1 - \alpha) : \alpha \in [0, 1] \}. \tag{5.5}$$

From Proposition 6.7 and Remark 6.8, we know that the convex non-increasing functions $\beta \mapsto g_b(Z_i, \beta)$ satisfy

$$\begin{cases} g_b(Z_i, \beta) = -\frac{Z_i^2}{4} \beta & \text{for } \beta \in \left[0, \frac{1}{2}\right] \\ g_b(Z_i, \beta) > -\frac{Z_i^2}{4} \beta & \text{for } \beta \in \left[\frac{1}{2}, 1\right]. \end{cases}$$

As a consequence for $Z_1 = Z_2$ we obtain that the minimum is uniquely attained for $\alpha = \frac{1}{2}$: we thus recover a more precise result than in Remark 4.6 where in that case any $\alpha \in [0, 1]$ would lead to a solution. Here, due to the correlation term C , each nucleus gets exactly one electron (see also Remark 5.11 below).

We now turn to the case $Z_1 > Z_2$. In that case it holds $g_b(Z_1, \beta) \leq g_b(Z_2, \beta)$ for all β . Moreover from the above properties of g_b it follows that the minimum in (5.5) is equal to

$$\min \left\{ g_b(Z_1, \alpha) - \frac{Z_2^2}{4} (1 - \alpha) : \alpha \in \left[\frac{1}{2}, 1 \right] \right\}.$$

Now, since $g_b(Z_1, 1) > -\frac{Z_1^2}{4}$ we may choose Z_2 close enough to Z_1 (yet keeping $Z_2 < Z_1$) such that

$$g_b(Z_1, 1) > -\frac{Z_1^2 + Z_2^2}{8},$$

in which case the minimum in (5.5) is attained for some $\alpha < 1$: this is quite different from what obtained in Remark 4.6 where the minimum would be only for $\alpha = 1$. Somehow this allows for a continuity of the solution set of (5.5) as Z_2 gets closer to Z_1 .

Since ρ represents the probability distribution of the N electrons, the presence of values of α_i which are not of the form k/N should be interpreted as the presence of a fractional number of electrons. This fact already appeared in the literature (see for instance [17,20]) and has a reasonable interpretation in terms of time-averaging.

Remark 5.11. We now discuss more extensively the H_2 molecule bond dissociation, and we show how our results compare to the results of [5, Theorem 5.1]. The H_2 molecule corresponds to $M = 2$ nuclei with charges $Z_1 = Z_2 = 1$ and $N = 2$ electrons. The physical total energy for this molecule when the nuclei are located at X_i/ε is given by

$$\frac{2\varepsilon}{|X_1 - X_2|} + \inf \{ 2T(\rho) + C(\rho) - 2(U_{X_1/\varepsilon}(\rho) + U_{X_2/\varepsilon}(\rho)) : \|\rho\| = 1 \}, \tag{5.6}$$

where the first term in the nucleus-nucleus Coulomb interaction and is vanishing with ε . The inf part corresponds to $2 \inf G_\varepsilon$ for $b = 1/2$. The representation of the Γ -limit of G_ε in this case is given by

$$G(\rho) = g_b(1, \rho(\{X_1\})) + g_b(1, \rho(\{X_2\}))$$

and, according to Remark 5.10 above, the minimum of G is attained for $\rho(\{X_1\}) = \rho(\{X_2\}) = \frac{1}{2}$. It follows that the energy (5.6) above converges, as $\varepsilon \rightarrow 0$ to

$$\begin{aligned} 4 g_b \left(1, \frac{1}{2} \right) &= 4 \min \left\{ T(\rho) - \mathcal{U}_0(\rho) : \|\rho\| = \frac{1}{2} \right\} \\ &= 2 \min \{ T(\rho) - U_0(\rho) : \|\rho\| = 1 \} \end{aligned}$$

which is twice the energy of an hydrogen atom as proved in [5, Theorem 5.1]. Note that with our normalization $\|\rho\| = \frac{1}{2}$ represents the energy of one electron when $N = 2$.

6. The general interacting case

In this section we consider the general case $b > 0$ and $N \geq 3$, for which the proof of existence as well as a full characterization of the Γ -limit G of the family of functionals $\{G_\varepsilon\}_\varepsilon$ seems a hard issue. In view of the successive steps of the preceding section Section 5 for the case $N = 2$, we expect to have the following properties on G over \mathcal{P}^- :

(P1) $G(\rho)$ only depends on the restriction of ρ to the nuclei X_1, \dots, X_M in the sense

$$G(\rho) = G(\rho^\# + \rho^\perp) = G(\rho^\#),$$

where we use the decomposition $\rho = \rho^\# + \rho^\perp$ of Section 5 with

$$\rho^\# = \sum_{i=1}^M \rho(\{X_i\})\delta_{X_i};$$

(P2) there exists a function g_b given as a generalization to any $N \geq 2$ of the Definition 5.2 such that

$$G(\rho^\#) = \sum_{i=1}^M g_b(Z_i, \rho(\{X_i\})).$$

Note that in the case $N = 2$ of Section 5, (P1) is obtained in Theorem 5.2 via the localization Lemma 5.1 which allows to control the transport term C . On the other hand, the proof of (P2) relies mainly on Lemma 5.6 which is linked to a deep understanding of the lower semicontinuous envelope \overline{C} of C , that is only fully characterized for this particular value of N .

In Subsection 6.1 below, we do obtain the properties (P1) and (P2) for the general case under some regularity assumptions on the subprobability ρ , which in particular is required to give small mass to the nuclei X_i (precisely $\rho(\{X_i\}) \leq 1/N$ for all i). Then in Subsection 6.2 we derive (P2) for the special case $M = 1$, *i.e.*, when there is only one nucleus.

6.1. Full characterization of G for particular subprobabilities

We consider the general case $b > 0$, $M \geq 1$ and $N \geq 2$.

We represent every measure $\rho \in \mathcal{P}^-$ as

$$\rho = \sum_{i=1}^M \alpha_i \delta_{X_i} + \rho^\perp$$

with ρ^\perp which does not charge the points X_i . In Proposition 3.7 we already showed that whenever $\alpha_1 = \dots = \alpha_M = 0$ then $G(\rho) = 0$. In this section we compute G in the case $\alpha_i \leq 1/N$ for all i . A particular case is $M = N$ and $\alpha_i = 1/N$ for all i which is the expected optimal configuration for the hydrogen molecule H_N .

In the following proof, we shall need the notion of concentration of a finite measure ρ , which is denoted $\mu(\rho)$ and defined by

$$\mu(\rho) := \lim_{r \rightarrow 0} \sup_x \rho(B(x, r)).$$

In particular $\mu(\alpha\delta_X) = \alpha$ and, in general $\mu(\rho) \geq \alpha$ implies that $\rho = \alpha\delta_X + \sigma$ for some point X and a nonnegative measure σ . We can now state our result.

Theorem 6.1. *Let $\rho = \alpha_1\delta_{X_1} + \dots + \alpha_M\delta_{X_M} + \rho^\perp$ with $\alpha_i \leq 1/N$ for all i and $\mu(\rho^\perp) \leq 1/N$, then*

$$G(\rho) = - \sum_{i=1}^M \alpha_i \frac{Z_i^2}{4}.$$

Proof. Since for every ρ

$$G^-(\rho) \geq G^0(\rho) = - \sum_{i=1}^M \alpha_i \frac{Z_i^2}{4},$$

it is enough to prove that

$$G^+(\rho) \leq - \sum_{i=1}^M \alpha_i \frac{Z_i^2}{4}.$$

We first consider the case $\rho \in \mathcal{P}$. By Theorem 4.4, there exists a sequence $\rho_\varepsilon \xrightarrow{*} \rho$ such that

$$G_\varepsilon^0(\rho_\varepsilon) = \varepsilon^2 T(\rho_\varepsilon) - \varepsilon U(\rho_\varepsilon) \longrightarrow - \sum_{i=1}^M \alpha_i \frac{Z_i^2}{4}.$$

Note that since $\rho \in \mathcal{P}$ the sequence $\{\rho_\varepsilon\}_\varepsilon$ converges narrowly to ρ and $\limsup \mu(\rho_\varepsilon) \leq \mu(\rho)$. Assume first that $\mu(\rho) < 1/N$. Then by a result of [6] (a proof of this result is also available in [24] or, in [3, Proposition 2.5] for the more restrictive case $\mu(\rho) < \frac{1}{N(N-1)^2}$) the transport cost C is uniformly bounded on the family $\{\rho_\varepsilon\}_\varepsilon$, so we have

$$\lim_{\varepsilon \rightarrow 0} \varepsilon C(\rho_\varepsilon) = 0.$$

We conclude that

$$G^+(\rho) \leq \limsup_{\varepsilon \rightarrow 0} G_\varepsilon(\rho_\varepsilon) = G^0(\rho) = - \sum_{i=1}^M \alpha_i \frac{Z_i^2}{4}.$$

We obtain the general case by approximation since G^+ is lower semicontinuous and the right-hand side is upper semicontinuous with respect to weak* convergence. \square

6.2. The special case $M = 1$ and $b > 0$

In this case, we may assume that the position of the unique nucleus is $X_1 = 0$. We prove (P2) with the following definition for the function g_b , which can be seen as the ground state energy of a molecule with only one nucleus.

Definition 6.2. For $R > 0, Z \in \mathbb{R}_+$ and $\alpha \in [0, 1]$ we define

$$g_b^R(Z, \alpha) := \inf \left\{ T(\rho) + bC(\rho) - Z U_0(\rho) : \rho \in \mathcal{P}, \int_{B(0,R)} d\rho \leq \alpha \right\}$$

and then set

$$g_b(Z, \alpha) := \sup \{ g_b^R(Z, \alpha) : R > 0 \}. \tag{6.1}$$

Note that the definition above could look ambiguous because we already defined the function g_b in (5.2). However, we shall obtain as a consequence of Theorem 6.4 that the two definitions of the function g_b given in (6.1) above and in (5.2) do coincide for the case $N = 2$ (see Remark 6.5 below).

Let us state some properties of the fonctions g_b and g_b^R that shall be useful below.

Lemma 6.3. For any $R > 0$, the functions $\alpha \mapsto g_b^R(Z, \alpha)$ is convex, continuous and non-increasing on $[0, 1]$ and $g_b^R(Z, \alpha) \geq -\frac{Z^2}{4}\alpha$. The same holds for $\alpha \mapsto g_b(Z, \alpha)$.

Proof. First note that the functions $g_b^R(Z, \cdot)$ and $g_b(Z, \cdot)$ may in fact be defined through the same formulas on $[0, +\infty[$, then being constant and equal to $g_b(Z, 1)$ on $[1, +\infty[$. Moreover, the functions $g_b^R(Z, \cdot)$ are clearly non-increasing, so that this also holds for $g_b(Z, \cdot)$. Finally the convexity of $\alpha \mapsto g_b^R(Z, \alpha)$ follows from the convexity of the map

$$\rho \mapsto G_1(\rho) = T(\rho) + bC(\rho) - Z U_0(\rho),$$

and taking the supremum over R yields the convexity for $g_b(Z, \cdot)$. It remains to show the continuity on $[0, 1]$. To see this we first compute

$$\forall R > 0, \forall \alpha \in [0, 1], \quad g_b^R(Z, \alpha) \geq \inf_{\rho \in \mathcal{P}} G_1^0(\rho) = \inf_{\rho \in \mathcal{P}} G^0 = -\frac{Z^2}{4},$$

where the last equality follows from (4.1). Consider now $R > 0$ and $\rho \in \mathcal{P}$ smooth and such that $\int_{B(0,R)} d\rho = 0$, then for all $\lambda > 1$ one has $\int_{B(0,R)} d\rho^{1/\lambda} = 0$ and

$$g_b^R(Z, 0) \leq G_1(\rho^{1/\lambda}) \leq \frac{1}{\lambda^2}T(\rho) + \frac{b}{\lambda}C(\rho) - \frac{Z}{\lambda}U_0(\rho)$$

and letting λ go to infinity yields $g_b^R(Z, 0) \leq 0$. From the preceding we thus get the continuity of $g_b^R(Z, \cdot)$ and $g_b(Z, \cdot)$ at $\alpha = 0$. Finally these convex non-increasing functions take their values in $\left[-\frac{Z^2}{4}, 0\right]$, so they are bounded and thus continuous on $]0, +\infty[$. □

We are now in position to prove our main result in this particular case.

Theorem 6.4. *For $M = 1$ and $X_1 = 0$ it holds*

$$\forall \alpha \in [0, 1], \quad G(\alpha\delta_0) = g_b(Z, \alpha) \tag{6.2}$$

where $g_b(Z, \alpha)$ is given by (6.1). Moreover, it holds

$$G^-(\alpha\delta_0 + \rho^\perp) \geq g_b(Z, \alpha) \tag{6.3}$$

for all $\alpha \in [0, 1]$ and for all ρ^\perp such that $\rho^\perp(\{0\}) = 0$ and $\alpha\delta_0 + \rho^\perp \in \mathcal{P}^-$.

Remark 6.5. In view of the above result, it follows from Theorem 5.7 applied to $\rho = \alpha\delta_0$ that the two definitions of g_b in (5.2) and (6.1) coincide in the case $N = 2$. Unfortunately, at the moment for $N \geq 3$ we do not have an explicit definition for g_b that would involve a kind of general partial transport as in (5.2).

Remark 6.6. It follows from (6.2) and (6.3) that, in the case $M = 1$, the minimum of the Γ -limit G is attained for any $\rho = \alpha\delta_0$ such that α minimizes $g_b(Z, \cdot)$ on $[0, 1]$: since this function is non-increasing, we note that in particular the probability δ_0 is thus a particular solution. It would be the unique solution in case $g_b(Z, \alpha)$ attains has unique minimum $\alpha = 1$ on $[0, 1]$, which seems a reasonable conjecture but still an open question.

Proof of Theorem 6.4. We first show (6.3). Take admissible α and ρ^\perp and consider a family $\{\rho_\varepsilon\}_{\varepsilon>0}$ in \mathcal{P} weakly* converging to $\alpha\delta_0 + \rho^\perp$. Fix $\eta > 0$, then for $r > 0$ small enough we have $\int_{B(0,r)} \rho^\perp \leq \eta/2$, so that for $\varepsilon > 0$ small enough it holds $\int_{B(0,r)} d\rho_\varepsilon \leq \alpha + \eta$. Thus denoting by $\rho_\varepsilon^\varepsilon(x)$ the rescaled version of ρ_ε

$$\rho_\varepsilon^\varepsilon(x) = \varepsilon^3 \rho_\varepsilon(\varepsilon x)$$

we have

$$\int_{B(0,r/\varepsilon)} d\rho_\varepsilon^\varepsilon \leq \alpha + \eta \quad \text{and} \quad G_\varepsilon(\rho_\varepsilon) = G_1(\rho_\varepsilon^\varepsilon) \geq g_b^{r/\varepsilon}(Z, \alpha + \eta)$$

where $g_b^R(Z, \cdot)$ is extended on \mathbb{R}_+ as in the proof of Lemma 6.3. This yields $\liminf_{\varepsilon \rightarrow 0} G_\varepsilon(\rho_\varepsilon) \geq g_b(Z, \alpha + \eta)$ and since this holds for any such family $\{\rho_\varepsilon\}_\varepsilon$ we infer $G^-(\alpha\delta_0 + \rho^\perp) \geq g_b(Z, \alpha + \eta)$. The claim then follows by continuity of $\alpha \mapsto g_b(Z, \alpha)$ on $[0, 1]$.

We now turn to (6.2), so we have to prove the inequalities

$$\forall \alpha \in [0, 1], \quad g_b(Z, \alpha) \leq G^-(\alpha\delta_0) \leq G^+(\alpha\delta_0) \leq g_b(Z, \alpha).$$

Note that $G^-(\alpha\delta_0) \geq g_b(Z, \alpha)$ follows from the preceding with $\rho^\perp = 0$. It remains to show $G^+(\alpha\delta_0) \leq g_b(Z, \alpha)$. We first note that this holds for $\alpha = 1$: indeed, in that case one has

$$g_b(Z, 1) = \inf\{G_1(\rho) : \rho \in \mathcal{P}\} \geq G^+(\delta_0),$$

where in the last inequality we again use that $G_1(\rho) = G_\varepsilon(\rho^{1/\varepsilon})$ for all $\varepsilon > 0$ and $\rho^{1/\varepsilon} \xrightarrow{*} \delta_0$ as $\varepsilon \rightarrow 0$. Let now α such that $0 \leq \alpha < 1$ and consider a family $\{\rho_R\}_{R>0}$ in \mathcal{P} such that

$$\int_{B(0,R)} d\rho_R \leq \alpha \quad \forall R > 0 \quad \text{and} \quad \lim_{R \rightarrow +\infty} G_1(\rho_R) = g_b(Z, \alpha).$$

Up to extracting a subfamily, we may assume that $\{\rho_R\}_{R>0}$ weakly* converges to some $\rho \in \mathcal{P}^-$. Then one has $\int d\rho = \beta$ for some $\beta \leq \alpha$. We infer:

- for fixed $\varepsilon > 0$, $\rho_R^{1/\varepsilon} \xrightarrow{*} \rho^{1/\varepsilon}$ as $R \rightarrow +\infty$,
- $\rho^{1/\varepsilon} \xrightarrow{*} \beta \delta_0$ as $\varepsilon \rightarrow 0$,

and since the weak* topology on \mathcal{P}^- is metrizable, we can extract a subfamily $\{\rho_{R_\varepsilon}\}_{\varepsilon>0}$ such that $\rho_{R_\varepsilon}^{1/\varepsilon} \xrightarrow{*} \beta \delta_0$ with $R_\varepsilon \rightarrow +\infty$. Now we compute

$$\limsup_{\varepsilon>0} G_\varepsilon(\rho_{R_\varepsilon}^{1/\varepsilon}) = \limsup_{\varepsilon>0} G_1(\rho_{R_\varepsilon}) = g_b(Z, \alpha).$$

Since $\rho_{R_\varepsilon}^\varepsilon \xrightarrow{*} \beta \delta_0$, this implies $G^+(\beta \delta_0) \leq g_b(Z, \alpha)$. If $\beta = \alpha$, the proof is complete. Otherwise $\beta < \alpha$ and thanks to the first step we infer

$$g_b(Z, \beta) \leq G^-(\beta \delta_0) \leq G^+(\beta \delta_0) \leq g_b(Z, \alpha).$$

Since $g_b(Z, \cdot)$ is convex non-increasing, this implies that this function is constant on $[\beta, 1]$. But then we have

$$G^+(\beta \delta_0) \leq g_b(Z, \alpha) = g_b(Z, 1) \quad \text{and} \quad G^+(\delta_0) \leq g_b(Z, 1) = g_b(Z, \alpha)$$

and by convexity of G^+ on $[\beta \delta_0, \delta_0]$ we get the desired inequality $G^+(\alpha \delta_0) \leq g_b(Z, \alpha)$. □

We conclude this subsection with some properties of the function $g_b(Z, \cdot)$.

Proposition 6.7. *It holds*

$$\forall \alpha \in \left[0, \frac{1}{N}\right], \quad g_b(Z, \alpha) = -\frac{Z^2}{4} \alpha$$

and

$$g_b(Z, \alpha) > -\frac{Z^2}{4} \alpha \quad \text{whenever } \overline{C}(\rho) > 0 \text{ for any } \rho \in \mathcal{P}^- \text{ such that } \int d\rho = \alpha.$$

Remark 6.8. In the case $N = 2$, it follows from Remark 2.6 that $\overline{C}(\rho) > 0$ for any $\rho \in \mathcal{P}^-$ such that $\int \rho > \frac{1}{2}$, so in that case $g_b(Z, \alpha) > -\frac{Z^2}{4} \alpha$ for any $\alpha > \frac{1}{2}$.

Proof of Proposition 6.7. It follows from Theorems 6.1 and 6.4 applied to $\rho = \alpha \delta_{X_1}$ that $g_b(Z, \alpha) = -\frac{Z^2}{4}\alpha$ whenever $\alpha \leq \frac{1}{N}$.

We now turn to the second claim, and assume that $\overline{C}(\rho) > 0$ for any $\rho \in \mathcal{P}^-$ such that $\int d\rho = \alpha$. From Lemma 6.3 we already know that $g_b(Z, \alpha) \geq -\frac{Z^2}{4}\alpha$, we assume by contradiction that $g_b(Z, \alpha) = -\frac{Z^2}{4}\alpha$. Then there exists a sequence $(\rho_n)_n$ in \mathcal{P} and a sequence $R_n \rightarrow +\infty$ such that

$$T(\rho_n) + bC(\rho_n) - Z U_0(\rho_n) \rightarrow -\frac{Z^2}{4}\alpha \quad \text{and} \quad \forall n, \int_{B(0,R_n)} d\rho_n \leq \alpha.$$

If we set $u_n := \sqrt{\rho_n}$ then $\int u_n^2 dx = 1$ for all n and since $C(\rho_n) \geq 0$ we get

$$\limsup_n \int \left[|\nabla u_n|^2 - Z \frac{u_n^2}{|x|} \right] dx = \limsup [T(\rho_n) - Z U_0(\rho_n)] \leq -\frac{Z^2}{4}\alpha.$$

Now up to extracting a subsequence we may assume that (u_n) weakly converges in $L^2(\mathbb{R}^3)$ to some function u and the above limsup is a limit, so we can apply Lemma 6.9 below and get

$$\int u^2 dx \geq \alpha \quad \text{and} \quad \int \left[|\nabla u|^2 - Z \frac{u^2}{|x|} \right] dx \leq \lim_n \int \left[|\nabla u_n|^2 - Z \frac{u_n^2}{|x|} \right] dx \leq -\frac{Z^2}{4}\alpha.$$

From the properties of ρ_n we infer that $\int_{B(0,R_n)} u_n^2 \leq \alpha$ for all n , so that $\int u^2 dx \leq \alpha$. This implies that $\int u^2 dx = \alpha$, and from (4.1) we get

$$\int \left[|\nabla u|^2 - Z \frac{u^2}{|x|} \right] dx \geq -\frac{Z^2}{4}\alpha.$$

Summarizing we obtain

$$\lim_n [T(\rho_n) - Z U_0(\rho_n)] = \lim_n \int \left[|\nabla u_n|^2 - Z \frac{u_n^2}{|x|} \right] dx = -\frac{Z^2}{4}\alpha$$

and then $\lim C(\rho_n) = 0$. On the other hand, $\liminf C(\rho_n) \geq \overline{C}(u^2) > 0$ since $\int u^2 = \alpha$, which is the desired contradiction. \square

Lemma 6.9. *Let (u_n) be a sequence in $H^1(\mathbb{R}^3)$ that weakly converges in $L^2(\mathbb{R}^3)$ to some function u , and such that*

$$\forall n, \int u_n^2 dx = 1, \quad \text{and} \quad \lim_n \int \left[|\nabla u_n|^2 - Z \frac{u_n^2}{|x|} \right] dx \leq -\frac{Z^2}{4}\alpha$$

for some $\alpha \in]0, 1]$. Then $(u_n)_n$ weakly converges in $H^1(\mathbb{R}^3)$ to u and

$$\int u^2 dx \geq \alpha \text{ and } \int \left[|\nabla u|^2 - Z \frac{u^2}{|x|} \right] dx \leq \lim_n \int \left[|\nabla u_n|^2 - Z \frac{u_n^2}{|x|} \right] dx \leq -\frac{Z^2}{4} \alpha.$$

Note that in the above result when $\alpha = 1$ it follows that $(u_n)_n$ strongly converges in $L^2(\mathbb{R}^3)$ to u and the function u^2 is a solution of the problem (4.1). Although all arguments for the proof can be found in [16], we provide them for the sake of completeness.

Proof. We first note that for n large enough one has

$$\int |\nabla u_n|^2 \leq Z \int \frac{u_n^2}{|x|} dx$$

which together with Lemma 3.6 yields that $(\nabla u_n)_n$ is bounded in $L^2(\mathbb{R}^3, \mathbb{R}^3)$. Since $(u_n)_n$ is also bounded in $L^2(\mathbb{R}^3)$, we infer that it is bounded in $H^1(\mathbb{R}^3)$, so it converges weakly in $H^1(\mathbb{R}^3)$ to u . By the weak lower semicontinuity of the H^1 seminorm we obtain

$$\int |\nabla u|^2 dx \leq \liminf_n \int |\nabla u_n|^2 dx.$$

Moreover, for every $R > 0$ we have

$$\limsup_n \int \frac{u_n^2}{|x|} dx \leq \limsup_n \left[\int_{B_R} \frac{u_n^2}{|x|} dx + \frac{1}{R} \right] = \int_{B_R} \frac{u^2}{|x|} dx + \frac{1}{R}$$

where B_R denotes the ball in \mathbb{R}^3 of radius R centered at the origin. Since R is arbitrary, we get

$$\limsup_n \int \frac{u_n^2}{|x|} dx \leq \int \frac{u^2}{|x|} dx.$$

Then we deduce

$$\int \left[|\nabla u|^2 - Z \frac{u^2}{|x|} \right] dx \leq \lim_n \int \left[|\nabla u_n|^2 - Z \frac{u_n^2}{|x|} \right] dx \leq -\frac{Z^2}{4} \alpha. \tag{6.4}$$

It remains to prove that $\int u^2 dx = \beta \geq \alpha$. Assume by contradiction that $\beta < \alpha$. We first note that $\beta > 0$ otherwise $u = 0$ which contradicts the inequalities in (6.4). Then the probability $\rho = \frac{1}{\beta} u^2$ satisfies

$$\int \left[\frac{|\nabla \rho|^2}{4\rho} - Z \frac{\rho}{|x|} \right] dx \leq -\frac{Z^2}{4} \frac{\alpha}{\beta}$$

which contradicts (4.1) since $\frac{\alpha}{\beta} > 1$. □

6.3. Perspectives and future work

A general proof (for any N and M) of (P1) and (P2) would give a full characterization of the Γ -limit functional G . It seems that a necessary tool for this program is an expression for the relaxation \overline{C} of the transport cost C with respect to the weak* convergence of measures. It would be then interesting to carry out a study (numerical and theoretical) of the minimizers of the Γ -limit functional G which could explain how the bond dissociation happens (*i.e.*, how the electrons are divided among the resulting molecules). A numerical study could also help to understand if the function $\alpha \mapsto g_b(Z, \alpha)$, which is non increasing in $[0, 1]$, attains its minimum uniquely for $\alpha = 1$ (see Remark 6.6).

Another interesting issue is the existence, for a fixed $\varepsilon > 0$, of minimizers $\rho_\varepsilon \in \mathcal{P}$ for the functional F_ε defined in (1.1). The existence of a solution $\bar{\rho}_\varepsilon \in \mathcal{P}^-$ for the relaxed functional \bar{F}_ε (with respect to the weak* convergence of measures) follows straightforwardly from the direct methods of the calculus of variations; the question if $\int d\bar{\rho}_\varepsilon = 1$ is sometimes called *ionization conjecture* and is part of our future work, together with a complete characterization of the relaxed correlation functional \overline{C} .

Appendix

A. Some notes on the model

The analysis of minimum problems for functionals involving the wave function is one of the most studied topics in quantum physics. The Born-Oppenheimer model for the electronic structure of several particles systems deals with the functional

$$\mathcal{E}(\psi) = \frac{\hbar^2}{2m} T(\psi) + U_{ee}(\psi) - U_{ne}(\psi), \quad (\text{A.1})$$

where

$$\begin{aligned} T(\psi) &= \sum_{s_i = \pm 1} \int_{\mathbb{R}^{3N}} \sum_{1 \leq i \leq N} |\nabla_{x_i} \psi|^2 dx_1 \dots dx_N \text{ (kinetic energy),} \\ U_{ee}(\psi) &= \sum_{s_i = \pm 1} \int_{\mathbb{R}^{3N}} \left(\sum_{1 \leq i < j \leq N} \frac{1}{|x_i - x_j|} \right) |\psi|^2 dx_1 \dots dx_N \text{ (electron-electron} \\ &\hspace{15em} \text{interaction),} \\ U_{ne}(\psi) &= \sum_{s_i = \pm 1} \int_{\mathbb{R}^{3N}} V(x_1, \dots, x_N) |\psi|^2 dx_1 \dots dx_N \text{ (electron-nuclei interaction).} \end{aligned}$$

We do not consider the nucleus-nucleus interaction because we assume that the nuclei are fixed and this extra term would then simply be a constant.

Here $\psi(x_1, s_1, x_2, s_2, \dots, x_N, s_N)$ is the wave function depending on space coordinates x_i and spin coordinates s_i , m the reduced mass of the nuclei, N the number of electrons, and V a potential. In the Coulomb case, if we assume to have M nuclei with positions X_k and charges Z_k ($k = 1, \dots, M$), we take

$$V(x_1, \dots, x_N) = \sum_{1 \leq k \leq M} \frac{Z_k}{|x - X_k|},$$

even if most of the analysis can be similarly carried out assuming only $V \in L^{3/2}(\mathbb{R}^3) + L^\infty(\mathbb{R}^3)$. The class where the functional above has to be minimized is the class of functions that, with respect to the space variables, belong to the Sobolev space $H^1(\mathbb{R}^{3N}; \mathbb{C})$ with $\|\psi\|_{L^2} = 1$.

Concerning the symmetry assumptions on the functions ψ in the admissible class, there are two main cases considered in physics:

- The *bosonic* case, where for all permutations σ of N points

$$\psi((x_{\sigma(1)}, s_{\sigma(1)}), \dots, (x_{\sigma(N)}, s_{\sigma(N)})) = \psi((x_1, s_1), \dots, (x_N, s_N));$$

- The *fermionic* case, where for all permutations σ of N points

$$\psi((x_{\sigma(1)}, s_{\sigma(1)}), \dots, (x_{\sigma(N)}, s_{\sigma(N)})) = \text{sign}(\sigma)\psi((x_1, s_1), \dots, (x_N, s_N)).$$

To simplify this rather complex problem, several approximated models have been proposed; as already said, here we deal with the one considered in the Density Functional Theory (DFT) introduced in the works of Thomas [25] and Fermi [10] and then revived by Hohenberg, Kohn and Sham [11, 12] and, from a variational point of view by Levy [14] and Lieb [15]. The DFT looks at the N -point distribution of electrons $\tilde{\rho}$ (also known as charge density) as the main variable, replacing the wave function ψ . The charge density $\tilde{\rho}$ is defined as follows, first define

$$\rho(x) = \sum_{s_i = \pm 1} \int_{\mathbb{R}^{3(N-1)}} |\psi(x, s_1, x_2, s_2, \dots, x_N, s_N)|^2 dx_2, \dots, dx_N,$$

and then $\tilde{\rho} = N\rho$. The potential (*i.e.*, electron-nuclei interaction) is easily expressed in terms of ρ or $\tilde{\rho}$ as

$$N \int V(x)\rho dx = \int V(x)\tilde{\rho} dx \quad (\text{potential}).$$

Again all the integrals with no domain of integration explicitly defined are intended on \mathbb{R}^3 . The kinetic energy cannot be expressed directly and possible approximations are given by the von Weizsäcker kinetic energy (which we considered in this paper)

$$\int \frac{|\nabla \tilde{\rho}|^2}{4\tilde{\rho}} dx$$

or by the Thomas-Fermi- von Weizsäcker kinetic energy

$$C_F \int \tilde{\rho}(x)^{5/3} dx + \int \frac{|\nabla \tilde{\rho}|^2}{4\tilde{\rho}} dx.$$

Both are local functionals and share the same scaling behavior with respect to the dilation operator $\rho \mapsto \rho^s := s^3 \rho(sx)$, which are the main properties used through this paper. The analysis could be quite different if one considers for instance the Kohn-Sham kinetic energy

$$F_{ks}(\rho) = \inf \left\{ \sum_{s_i = \pm 1} \int_{\mathbb{R}^{3N}} \sum_{1 \leq i \leq N} |\nabla_{x_i} \psi|^2 dx_1 \dots dx_N, : \right. \\ \left. \rho(x) = \sum_{s_i = \pm 1} \int_{\mathbb{R}^{3(N-1)}} |\psi(x, s_1, x_2, s_2, \dots, x_N, s_N)|^2 dx_2, \dots, dx_N \right\},$$

where the inf is taken on the set of Bosonic or Fermionic wave functions depending on the case at hand. The functional F_{ks} still has the same behavior with respect to the dilation operator above but has not in general an integral expression. In fact, as proved in [15, Theorems 1.1 and 1.2], the von Weizsäcker energy, up to a multiplicative constant, coincides with the F_{ks} energy in the Bosonic case while it only controls it from above and from below in the Fermionic case.

Concerning the approximation for electron-electron correlation, we have expressed it using the multi-marginal mass transport theory. When the Coulomb correlation term C above is chosen, the theory is usually called Strongly Correlated Electrons Density Functional Theory (SCE-DFT); it was started and developed since the late 90's (see for instance [21–23]) and the connection with optimal transport was made in [4, 7].

The problem of bond dissociating limit in the Born-Oppenheimer model has been considered in [19] where it has been proved that *as the nuclei of a molecule are dissociated into spatially separated clusters, the discrete molecular energies approach sums of the energies of isolated subsystems*. Although it is not clear that, in such a model, a neutral molecule dissociates into neutral subsystems, each of the subsystems obtained in [19] contains an integer number of electrons. This is not the case when using most of the current models in DFT.

References

- [1] H. ATTOUCH, “Variational Convergence for Functions and Operator”, Applicable Mathematics Series, Pitman (Advanced Publishing Program), Boston, 1984.
- [2] U. BINDINI and L. DEPASCALE, *Optimal transport with Coulomb cost and the semiclassical limit of density functional theory*, J. Éc. Polytech. Math. **4** (2017), 909–934.

- [3] G. BUTTAZZO, T. CHAMPION and L. DE PASCALE, *Continuity and estimates for multimarginal optimal transportation problems with singular costs*, Appl. Math. Optim. **78** (2018), 185–200.
- [4] G. BUTTAZZO, L. DE PASCALE and P. GORI-GIORGI, *Optimal-transport formulation of electronic density-functional theory*, Phys. Rev. A **85** (2012), 062502, 1–11.
- [5] H. CHEN, G. FRIESECKE and C. B. MENDEL, *Numerical methods for a Kohn-Sham density functional model based on optimal transport*, J. Chem. Theory Comput. **10** (2014), 4360–4368.
- [6] M. COLOMBO, S. DI MARINO and F. STRA, *Continuity of multi-marginal optimal transport with repulsive cost*, SIAM J. Math. Anal. **51** (2019), 2903–2926.
- [7] C. COTAR, G. FRIESECKE and C. KLÜPPELBERG, *Density functional theory and optimal transportation with Coulomb cost*, Comm. Pure Appl. Math. **66** (2013), 548–599.
- [8] C. COTAR, G. FRIESECKE and C. KLÜPPELBERG, *Smoothing of transport plans with fixed marginals and rigorous semiclassical limit of the Hohenberg-Kohn functional*, Arch. Ration. Mech. Anal. **228** (2018), 891–922.
- [9] G. DAL MASO, “An Introduction to Γ -Convergence”, Progress in Nonlinear Differential Equations and their Applications (PNLDE), Vol. 8, Birkhäuser-Verlag, Basel, 1993.
- [10] E. FERMI, *Un metodo statistico per la determinazione di alcune proprietà dell’atomo*, Rend. Accad. Naz. Lincei **6** (1927), 602–607.
- [11] P. HOHENBERG and W. KOHN, *Inhomogeneous electron gas*, Phys. Rev. B **136** (1964), 864–871.
- [12] W. KOHN and L. J. SHAM, *Self-consistent equations including exchange and correlation effects*, Phys. Rev. A **140** (1965), 1133–1138.
- [13] M. LEWIN, *Semi-classical limit of the Levy-Lieb functional in Density Functional Theory*, C. R. Math. Acad. Sci. Paris **356** (4) (2018), 449–455.
- [14] M. LEVY, *Universal variational functionals of electron densities, first-order density matrices, and natural spin-orbitals and solution of the v -representability problem*, Proc. Nat. Acad. Sci. U.S.A **76** (1979), 6062–6065.
- [15] E. H. LIEB, *Density functionals for Coulomb systems*, Internat. J. Quantum Chem. **24** (3) (1983), 243–277.
- [16] E. H. LIEB and M. LOSS, “Analysis”, Graduate Studies in Mathematics, Vol. 14, American Mathematical Society, Providence, 1997.
- [17] A. MIRTSCHINK, M. SEIDL and P. GORI-GIORGI, *Derivative discontinuity in the strong-interaction limit of density functional theory*, Phys. Rev. Lett. **111** (2013), 126402.
- [18] A. MIRTSCHINK, C. J. UMRIGAR, J. D. MORGAN III and P. GORI-GIORGI, *Energy density functionals from the strong-coupling limit applied to the anions of the He isoelectronic series*, J. Chem. Phys. **140** (2014), 18A532.
- [19] J. D. MORGAN and B. SIMON, *Behavior of molecular potential energy curves for large nuclear separations*, Internat. J. Quantum Chem. **17** (1980), 1143–1166.
- [20] J. P. PERDEW, R. G. PARR, M. LEVY and J. L. BALDUZ JR., *Density functional theory for fractional particle number: derivative discontinuity of the energy*, Phys. Rev. Lett. **49** (1982).
- [21] M. SEIDL, *Strong-interaction limit of density-functional theory*, Phys. Rev. A **60** (6) (1999), 4387–4395.
- [22] M. SEIDL, P. GORI-GIORGI and A. SAVIN, *Strictly correlated electrons in density-functional theory: A general formulation with applications to spherical densities*, Phys. Rev. A **75** (4) (2007), 042511.
- [23] M. SEIDL, J. P. PERDEW and M. LEVY, *Strictly correlated electrons in density-functional theory*, Physical Review A **59** (1) (1999), 51–54.
- [24] F. STRA, “Classical and Multi-marginal Optimal Transport, with Applications”, PhD Thesis, Sc. Norm. Super., Pisa, 2018.

- [25] L. H. THOMAS, *The calculation of atomic fields*, Math. Proc. Cambridge Philos. Soc. **23** (1927), 542–548.

Laboratoire IMATH
Université de Toulon
BP 20132
83957 La Garde Cedex, France
bouchitte@univ-tln.fr

Dipartimento di Matematica
Università di Pisa
Largo B. Pontecorvo, 5
56127 Pisa, Italia
giuseppe.buttazzo@unipi.it

Laboratoire IMATH
Université de Toulon
BP 20132
83957 La Garde Cedex, France
champion@univ-tln.fr

Dipartimento di Matematica e Informatica
Università di Firenze
Viale Morgagni 67/a
50134 Firenze, Italia
luigi.depascale@unifi.it